

# Self-introduction



My name is Atsumu Suzuki  
(鈴木州) from Kobe University  
Member of SK, HK, and T2K  
(Neutrino experiments)



Scuba diving  
(a couple of times a year)



Tennis  
weekend



Skating in winter



A photograph of three astronauts in space, floating against the backdrop of the Earth. The astronauts are wearing white space suits and are holding onto a structure. The Earth's surface is visible, showing blue oceans and brown landmasses. The title 'Introduction to Cosmic Ray' is overlaid in yellow text.


# Introduction to Cosmic Ray

**Atsumu Suzuki**  
Kobe University

# What are Cosmic Rays ?

- **Particles from the outside of the Earth (1ry cosmic rays)**
  - 90% of them are hydrogen nuclei (protons)
  - They interact with nitrogen and oxygen nuclei in the atmosphere and generate **2ry cosmic rays**.
    - **Muons** and **neutrinos** are also generated.
- **Typical CR muon energy is  $\sim 1-100$  GeV**

# Cosmic Ray Spectrum



# Why $10^{20} \sim 10^{21}$ eV ?

High energy protons interact with Cosmic Microwave Background (CMB):



What is the threshold energy  $E_p$  of such a proton ?

$$(E_p + E_{\text{CMB}})^2 - (p_p - E_{\text{CMB}})^2 = (m_n + m_\pi)^2$$

Assuming  $E_p = p_p$  because of high energy, we obtain

$$E_p = \frac{(m_n + m_\pi)^2}{4E_{\text{CMB}}}$$

$$E_{\text{CMB}} = 8.62 \times 10^{-5} [\text{eV K}^{-1}] \times 2.7 [\text{K}] = 2.33 \times 10^{-4} [\text{eV}],$$

$$\text{and } m_n = 940 [\text{MeV}], m_\pi = 140 [\text{MeV}] \rightarrow E_p = 1.25 \times 10^{21} \text{ eV}$$

(consistent with measurements)

The number of UHE CR coming to the Earth is suddenly suppressed at the energy of  $\sim 10^{20}$  eV.  $\Rightarrow$

**Greisen-Zatsepin-Kuzmin (GZK) Cutoff**



# Primary cosmic-ray flux

## Chemical composition





Figure 30.1: Fluxes of nuclei of the primary cosmic radiation in particles per energy-per-nucleus are plotted vs energy-per-nucleus using data from Refs. [1–15]. The inset shows the H/He ratio as a function of rigidity [1,3].

## 1ry CR proton energy distribution




# Balloon-borne Experiment with a Superconducting Spectrometer (BESS)




BESS-2000

- Collaboration between Japan and US (KEK, Univ. of Tokyo, Kobe Univ., JAXA, NASA, and Univ. of Maryland)
- **Purposes**
  1. Precision measurement of low energy 1ry CR antiprotons
  2. Search for CR antimatter (anti He nucleus)
  3. Precision measurement of 1ry CR proton and helium energy spectrums etc.
- Site: Lynn Lake (Canada), Antarctica




# Alpha Magnetic Spectrometer (AMS)

- Particle physics detector on the international space station for the cosmic ray measurement.



- Large area and solid angle
- Superconducting magnet + Si detector
- Good particle identification (PID)
  - TRD
  - RICH
  - ECAL  $15X_0$
- Total weight: 6t





# Cosmic ray flux measurements

**Japanese American Cooperative Emulsion Experiment (JACEE)** : Direct measurements of 1ry CR components and energy spectrum in Antarctica Balloon-borne experiment

## **Russia-Nippon Joint Balloon Experiment (RUNJOB)**

purpose: measuring the chemical compositions and energy spectra of the primary cosmic ray, balloon, Russia

## **HEAT (High-Energy Antimatter Telescope)**

purpose: study of CR  $e^-e^+$ , isotopic composition, balloon, New Mexico & Lynn Lake (Canada)

## **TRACER (Transition Radiation Array for Cosmic Energetic Radiation)**

purpose: direct measurements of the heavier primary cosmic-ray nuclei at high energies, balloon, Antarctica

## **ATIC (Advanced Thin Ionization Calorimeter)**


purpose: measuring the energy and composition of cosmic rays, balloon, Antarctica

## **CREAM (Cosmic Ray Energetics and Mass)**


purpose: determining the composition of cosmic rays up to the  $10^{15}$  eV (also known as the "knee prospect") in the cosmic ray spectrum, balloon, Antarctica

# Why at high latitude ?

To lower the cut-off rigidity  $R_c$ . The rigidity  $R = p/z$ , where  $p$  is the momentum and  $z$  is the charge ( $R$  for a proton of  $p = 1$  [GeV] is 1 [GV], and for a helium of  $p = 1$  [MeV] is 0.5 [MV]).



Plan view from the north pole




Estimate  $R_c$ . Magnetic field  $B \approx 5 \times 10^{-5}$  [T] at the altitude  $h \leq \sim 1,000$  km.

$$R_c = 0.3hB$$


$$= 0.3 \times 10^6 \times 5 \times 10^{-5} = 15 \text{ [GV]}$$

Tokyo:  $R_c \approx 11$  GV

# Why especially in Antarctica ?



One way track  
~ 20 ~ 30 hour flight




Circling orbit  
~1 month flight




# East-West effect

Charged cosmic rays receive a Lorentz force from the geomagnetic field. Due to the direction of the field from the south to north, positively charged particles from the west receive the force to outside and those from the east receive the force to inside. Therefore we observe the particle from the west more than those from the east.



# Result of SK

- $\nu$ 's from west  $>$   $\nu$ 's from east  
→ more positively charged particles than negatively charged ones
- Agreement between the data and simulation → correctness of our understanding about atmospheric neutrinos



Points : data

Boxes : simulation

PhysRevD.94.052001

# 2ry cosmic ray generation

1ry cosmic rays interact with N, O, C, etc. in the atmosphere.


Primary cosmic ray

p, He ...

atmosphere

Production height

10~30km



p-A cross-section ( 10 ~ 20GeV )  
 $\sim (40\text{mb})A^{2/3}$

mean free path

$\sim 40A^{-1/3} \text{ g}\cdot\text{cm}^{-2}$

→ Typical altitude of  
the 1<sup>st</sup> interaction  
 $\sim 15\text{km}$

\* 1 b (barn) =  $10^{-28} \text{ m}^2$



# 2ry cosmic ray generation


1ry cosmic rays interact with N, O, C, etc. in the atmosphere.

Primary cosmic ray

p, He ...

atmosphere

Generation height  
10~30km



2) Decay of  $\pi$

How far can 5 GeV charged pions travel ?

$$\gamma c \tau_\pi = \frac{E_\pi}{m_\pi} c \tau_\pi = 278 \text{ m} \sim 300 \text{ m}$$

(3.6 g/cm<sup>2</sup>)

\*note: interaction length of  $\pi \sim 160 \text{ g/cm}^2$

Most pions decay

$$\pi \rightarrow \mu + \nu_\mu$$

$$c = 3 \times 10^8 \text{ m/s}, m_\pi = 140 \text{ MeV}, \tau_\pi = 2.6 \times 10^{-8} \text{ sec}$$

# 2ry cosmic ray generation


1ry cosmic rays interact with N, O, C, etc. in the atmosphere.

Primary cosmic ray

p, He ...

atmosphere

Generation height  
10~30km



3) fate of  $\mu$

2.5 GeV muons

from life ~15 km

energy loss  $\sim 2\text{GeV}$

$\left\{ \begin{array}{l} \text{decay } \mu \rightarrow e + \nu_{\mu} + \nu_e \\ \text{hit the ground} \\ \text{absorption } (\mu^-) \\ \text{decay} \end{array} \right.$

Lower energy  $\mu$

Decay

Higher energy  $\mu$

Can not decay

$$\frac{v_{\mu}}{v_e} \sim 2 \quad (< \sim 1 \text{ GeV})$$

$$\frac{v_{\mu}}{v_e} > 2 \quad (> \sim 1 \text{ GeV})$$

SK

# Zenith angle and energy distributions of cosmic ray

Since the energy loss is  $2 \text{ MeV}/(\text{g}/\text{cm}^2)$  for high energy charged particles, minimum energy  $E_{min}$  of the particle which can reach the surface is


$$E_{min} = \frac{2 [\text{GeV}]}{\cos \theta}$$

Number of incident particles:

$$\begin{aligned} N(E > E_{min}) &= \int_{E_{min}}^{\infty} n(E) dE \\ &\propto \int_{E_{min}}^{\infty} E^{-\gamma} dE \\ &= \frac{2^{1-\gamma}}{\gamma-1} \cos^{\gamma-1} \theta \end{aligned}$$

• From measurements, we can get

$$\gamma - 1 \approx 2 \rightarrow \gamma \approx 3 \text{ (2.7)}$$







# Simple experiment (1)

## (CR flux measurement)

Measurement of the zenith angle distribution of **cosmic rays** by taking a **coincidence** of a pair of **scintillation counters**.




### Function of a discriminator



# Simple cosmic ray experiment(1)

Measurement of the zenith angle distribution of **cosmic rays** by taking a **coincidence** of a pair of **scintillation counters**.



**Discriminator**

outputs a pulse signal when an input signal is higher than the threshold.

**Coincidence module**

**Scaler**

counts the number of input pulses.

**Discriminator**

takes a coincidence between input signals

$L$ : distance between the scintillators

$S$ : area of the scintillator

$N$ : number of events

$T$ : measurement time


$\Omega$ : solid angle

Flux:  $\Phi = \frac{N}{S\Omega T} \approx \frac{L^2 N}{S^2 T} \sim 100 \text{ m}^{-2}\text{s}^{-1}\text{sr}^{-1}$  (typical sea-level value about vertical direction)

$\propto \cos^2 \theta \rightarrow \gamma - 1 \approx 2 \rightarrow \gamma \approx 3 \text{ (2.7)}$  :consistent with the measurements




# Simple experiment (2)

(muon lifetime measurement)



# Simple experiment (2)

(muon lifetime measurement)



# Simple experiment (2)

## Exercise (homework)

1. Particle decay follows the following formula:

$$N = N_0 e^{-t/\tau},$$

where  $N$  and  $N_0$  are the numbers of events at time  $t$  and 0, respectively, and  $\tau$  is the lifetime.

The right side table is a result of the muon decay experiment.

(a) Make a plot of the number of events as a function of the decay time. Use log scale as a vertical axis.

(b) Get the muon lifetime.

2. Some events show no 2<sup>nd</sup> signal which corresponds to decay electron. What are those events ?


\* Note that the condition in the next page.

Decay time $t$ [ $\mu$ sec]	Number of events	Decay time $t$ [ $\mu$ sec]	Number of events
0.3	1501	4.7	194
0.5	1308	4.9	189
0.7	1191	5.1	155
0.9	1082	5.3	157
1.1	1024	5.5	127
1.3	886	5.7	134
1.5	823	5.9	102
1.7	775	6.1	90
1.9	700	6.3	96
2.1	610	6.5	79
2.3	544	6.7	62
2.5	547	6.9	85
2.7	497	7.1	65
2.9	463	7.3	61
3.1	422	7.5	53
3.3	380	7.7	66
3.5	340	7.9	41
3.7	291	8.1	28
3.9	276	8.3	29
4.1	283	8.5	26
4.3	245	8.7	27
4.5	204		

\* ex. "0.3" means  $0.1 \leq t < 0.4$  [ $\mu$  sec]




# Real Geometry of the Detector



Since S2 is small and the distance between S1 and S3 is long enough, particles which pass through S1, S2, and S3 always reach S4.

(There is no event which passes through S1, S2, and S3, but does not pass S4 like the following one: )



# Summary

## **As introduction to cosmic rays :**

- **What are cosmic rays ?**
- **Cosmic ray spectrum**
- **Geomagnetic effect**
- **2ry CR generation – including atmospheric neutrinos –**
- **Simple experiments – Student experiments of Kobe Univ.**

**In addition to above, I gave you a homework (it is not a duty but I would like you to do).**



Thank you (^o^)



Son-san