

Self-introduction



My name is Atsumu Suzuki
(鈴木州) from Kobe
University
Member of SK, HK, and T2K



Scuba diving
(a couple of times a year)



Tennis
weekend



My shoes

Skating in winter

A photograph of astronauts in space, likely on the International Space Station, with the Earth's horizon and atmosphere visible in the background. The astronauts are wearing white suits and are positioned in the foreground, with their bodies and limbs visible against the dark space and the blue and white of the Earth. The Earth's surface shows a mix of blue oceans and brownish-green landmasses.


Introduction to Cosmic Ray

Atsumu Suzuki
Kobe University

What are Cosmic Rays ?

- **Particles from the outside of the Earth (1ry cosmic rays)**
 - 90% of them are hydrogen nuclei (protons)
 - They interact with nitrogen and oxygen nuclei in the atmosphere and generate **2ry cosmic rays**.
 - **Muons** and **neutrinos** are also generated.
- **Typical CR muon energy is $\sim 1-100$ GeV**

Cosmic Ray Spectrum



Why $10^{20} \sim 10^{21}$ eV ?

High energy protons interact with Cosmic Microwave Background (CMB):



What is the threshold energy E_p of such a proton ?

$$(E_p + E_{\text{CMB}})^2 - (p_p - E_{\text{CMB}})^2 = (m_n + m_\pi)^2$$

Assuming $E_p = p_p$ because of high energy, we obtain

$$E_p = \frac{(m_n + m_\pi)^2}{4E_{\text{CMB}}}$$

$$E_{\text{CMB}} = 8.62 \times 10^{-5} [\text{eV K}^{-1}] \times 2.7 [\text{K}] = 2.33 \times 10^{-4} [\text{eV}],$$

$$\text{and } m_n = 940 [\text{MeV}], m_\pi = 140 [\text{MeV}] \rightarrow E_p = 1.25 \times 10^{21} \text{ eV}$$

(consistent with measurements)

The number of UHE CR coming to the Earth is suddenly suppressed at the energy of $\sim 10^{20}$ eV. \Rightarrow

Greisen-Zatsepin-Kuzmin (GZK) Cutoff

Primary cosmic-ray flux

Chemical composition





Figure 30.1: Fluxes of nuclei of the primary cosmic radiation in particles per energy-per-nucleus are plotted vs energy-per-nucleus using data from Refs. [1–15]. The inset shows the H/He ratio as a function of rigidity [1, 3].

1ry CR proton energy distribution




Balloon-borne Experiment with a Superconducting Spectrometer (BESS)




BESS-2000

- Collaboration between Japan and US (KEK, Univ. of Tokyo, Kobe Univ., JAXA, NASA, and Univ. of Maryland)
- **Purposes**
 1. Precision measurement of low energy 1ry CR antiprotons
 2. Search for CR antimatter (anti He nucleus)
 3. Precision measurement of 1ry CR proton and helium energy spectrums etc.
- Site: Lynn Lake (Canada), Antarctica




Alpha Magnetic Spectrometer (AMS)

- Particle physics detector on the international space station for the cosmic ray measurement.



- Large area and solid angle
- Superconducting magnet + Si detector
- Good particle identification (PID)
 - TRD
 - RICH
 - ECAL $15X_0$
- Total weight: 6t



Cosmic ray flux measurements

Japanese American Cooperative Emulsion Experiment (JACEE) : Direct measurements of 1ry CR components and energy spectrum in Antarctica Balloon-borne experiment

Russia-Nippon Joint Balloon Experiment (RUNJOB)

purpose: measuring the chemical compositions and energy spectra of the primary cosmic ray, balloon, Russia

HEAT (High-Energy Antimatter Telescope)

purpose: study of CR e^-e^+ , isotopic composition, balloon, New Mexico & Lynn Lake (Canada)

TRACER (Transition Radiation Array for Cosmic Energetic Radiation)

purpose: direct measurements of the heavier primary cosmic-ray nuclei at high energies, balloon, Antarctica

ATIC (Advanced Thin Ionization Calorimeter)


purpose: measuring the energy and composition of cosmic rays, balloon, Antarctica

CREAM (Cosmic Ray Energetics and Mass)


purpose: determining the composition of cosmic rays up to the 10^{15} eV (also known as the "knee prospect") in the cosmic ray spectrum, balloon, Antarctica

Why at high latitude ?

To lower the cut-off rigidity R_c . The rigidity $R = p/z$, where p is the momentum and z is the charge (R for a proton of $p = 1$ [GeV] is 1 [GV], and for a helium of $p = 1$ [MeV] is 0.5 [MV]).



Plan view from the north pole




Estimate R_c . Magnetic field $B \approx 5 \times 10^{-5}$ [T] at the altitude $\leq \sim 1,000$ km.

$$R_c = 0.3hB$$


$$= 0.3 \times 10^6 \times 5 \times 10^{-5} = 15 \text{ [GV]}$$

Tokyo: $R_c \approx 11$ GV

Why especially in Antarctica ?




One way track
~ 20 ~ 30 hour flight



Circling orbit
~1 month flight

East-West effect

Charged cosmic rays receive a Lorentz force from the geomagnetic field. Due to the direction of the field from the south to north, positive particles from the west receive the force to outside and those from the east receive the force to inside. Therefore we observe the particle from the west more than those from the east.



2ry cosmic ray generation

1ry cosmic rays interact with N, O, C, etc. in the atmosphere.


Primary cosmic ray

p, He ...

atmosphere

Production height

10~30km



p-A cross-section (10 ~ 20GeV)

$$\sim (40\text{mb})A^{2/3}$$

mean free path

$$\sim 40A^{-1/3} \text{ g}\cdot\text{cm}^{-2}$$

→ Typical altitude of the 1st interaction
~ 15km

* 1 b (barn) = 10^{-28} m^2

2ry cosmic ray generation


1ry cosmic rays interact with N, O, C, etc. in the atmosphere.

Primary cosmic ray

p, He ...

atmosphere

Generation height
10~30km



SK

2) Decay of π

How far can 5 GeV charged pions travel ?

$$\gamma c \tau_{\pi} = \frac{E_{\pi}}{m_{\pi}} c \tau_{\pi} = 278 \text{ m} \sim 300 \text{ m}$$

(3.6 g/cm²)

*note: interaction length of $\pi \sim 160 \text{ g/cm}^2$

Most pions decay

$$\pi \rightarrow \mu + \nu_{\mu}$$

$$c = 3 \times 10^8 \text{ m/s}, m_{\pi} = 140 \text{ MeV}, \tau_{\pi} = 2.6 \times 10^{-8} \text{ sec}$$

2ry cosmic ray generation


1ry cosmic rays interact with N, O, C, etc. in the atmosphere.

Primary cosmic ray

p, He ...

atmosphere

Generation height
10~30km



3) fate of μ

2.5 GeV muons

from life ~15 km

energy loss $\sim 2\text{GeV}$

$\left\{ \begin{array}{l} \text{decay } \mu \rightarrow e + \nu_{\mu} + \nu_e \\ \text{hit the ground} \\ \text{absorption } (\mu^-) \\ \text{decay} \end{array} \right.$

Lower energy μ

Decay

Higher energy μ

Can not decay

$$\frac{v_{\mu}}{v_e} \sim 2 \quad (< \sim 1 \text{ GeV})$$

$$\frac{v_{\mu}}{v_e} > 2 \quad (> \sim 1 \text{ GeV})$$

SK


Zenith angle and energy distributions of cosmic ray

Since the energy loss is $2 \text{ MeV}/(\text{g}/\text{cm}^2)$ for high energy charged particles, minimum energy E_{min} of the particle which can reach the surface is

$$E_{min} = \frac{2 [\text{GeV}]}{\cos \theta}$$

Number of incident particles:


$$\begin{aligned} N(E > E_{min}) &= \int_{E_{min}}^{\infty} n(E) dE \\ &\propto \int_{E_{min}}^{\infty} E^{-\gamma} dE \\ &= \frac{2}{\gamma - 1} \cos^{\gamma-1} \theta \end{aligned}$$



Simple experiment (1)


(CR flux measurement)

Measurement of the zenith angle distribution of **cosmic rays** by taking a **coincidence** of a pair of **scintillation counters**.



Simple cosmic ray experiment(1)

Measurement of the zenith angle distribution of **cosmic rays** by taking a **coincidence** of a pair of **scintillation counters**.



Discriminator

outputs a pulse signal when an input signal is higher than the threshold.

Coincidence module

Scaler

counts the number of input pulses.

Discriminator

takes a coincidence between input signals

L : distance between the scintillators

S : area of the scintillator

N : number of events

T : measurement time


Ω : solid angle

Flux: $\Phi = \frac{N}{S\Omega T} \approx \frac{L^2 N}{S^2 T} \sim 100 \text{ m}^{-2}\text{s}^{-1}\text{sr}^{-1}$ (typical sea-level value about vertical direction)

$\propto \cos^2 \theta \rightarrow \gamma - 1 \approx 2 \rightarrow \gamma \approx 3$ (2.7) :consistent with the measurements




Simple experiment (2)

(muon lifetime measurement)



Simple experiment (2)

(muon lifetime measurement)



Simple experiment (2)

Exercise

1. Particle decay follows the following formula:

$$N = N_0 e^{-t/\tau},$$

where N and N_0 are the numbers of events at time t and 0, respectively, and τ is the lifetime.

The right side table is a result of the experiment.


(a) Make a plot of the number of events as a function of the decay time (use log scale as a vertical axis).

(b) Get the muon lifetime.

2. Some events show no 2nd signal which corresponds to decay electron. What are those events ?

Decay time t [μ sec]	Number of events	Decay time t [μ sec]	Number of events
0.3	1501	4.7	194
0.5	1308	4.9	189
0.7	1191	5.1	155
0.9	1082	5.3	157
1.1	1024	5.5	127
1.3	886	5.7	134
1.5	823	5.9	102
1.7	775	6.1	90
1.9	700	6.3	96
2.1	610	6.5	79
2.3	544	6.7	62
2.5	547	6.9	85
2.7	497	7.1	65
2.9	463	7.3	61
3.1	422	7.5	53
3.3	380	7.7	66
3.5	340	7.9	41
3.7	291	8.1	28
3.9	276	8.3	29
4.1	283	8.5	26
4.3	245	8.7	27
4.5	204		

* ex. "0.3" means $0.1 \leq t < 0.4$ [μ sec]



Thank you (^o^)

