



UCL

MINOS: Unpacking the NuMI Beam Simulation Using the Near Detector

Alex Sells




MINOS

Background: Gustave Doré

MINOS detector

- Long baseline 735km neutrino oscillation experiment
- Looked for disappearance of ν_{μ} at far detector to measure oscillation parameters
- Active from 2005 to 2016 and in its day made the most precise Δm^2_{32} and $\sin^2\theta_{23}$ measurements



NuMI Beam

Hadron Production

Focusing/optics

~1km

Detection

120GeV
Protons

Target coordinate
reference $z=0\text{cm}$

Magnetic focusing horns

Graphite Target

Current $0 < I < 200 \text{ kA}$

Parent hadron –
predominantly pions

Hadron momentum
coordinates

p_z
 p_T (radial)

Predominantly
muons + muon
neutrinos from pion
decay



MINOS Near Detector

Calorimetry + charge
measurement

Why are we still interested in MINOS





- MINOS took lots of data in many different beam configurations, providing neutrino energy spectra for a range of conditions
- Near Detector data allows us to make inferences about the NuMI beam simulation, which is relevant to contemporary experiments' beam design



Simulation of the NuMI beam



- MC model of the ND energy spectrum using the G4NuMI-FLUGG simulation have imperfect agreement with the data
- A study identified that a major source of error was the hadron production in the NuMI target due to disagreement of up to 30% across different theoretical models




Model	$\langle p_T \rangle$ (GeV/c)
Fluka 2001[23]	0.43
Fluka 2005[23]	0.36
MARS-v.15[17]	0.38
Malensek[22]	0.50
Geant/Fluka[19]	0.37
Sanford-Wang[20]	0.42
CKP[21]	0.44

Mean transverse momenta of π^+ from a graphite target struck by 120GeV protons, as calculated by several interaction Monte Carlo's. Taken from [1]

[1] S. Kopp, Z. Pavlovic, P. Vahle and R. Ospanov, Constraining the Beam Neutrino MC Flux Using the MINOS ND Data, MINOS Doc-DB #2965.

Hadron Production


Projections of x_T bins along p_T axis: had. prod. MC and two parameterisations



- FLUGG prediction for hadron production momentum
- Reweight this by fitting to data
- Requires good parametrization

Hadron Production

Projections of x_F bins along p_T axis: had. prod. MC and two parameterisations




$$\frac{d^2 N}{dx_F dp_T} = [A(x_F) + B(x_F)p_T] \exp(-C(x_F)p_T^{\frac{2}{3}})$$

- Earlier attempt –standard BMPT parametrization
- Combines theoretical and empirical terms
- Fit to the different MC x_F bands but imperfect model

Hadron Production

Projections of x_F bins along p_T axis: had. prod. MC and two parameterisations







$$\frac{d^2 N}{dx_F dp_T} = (B(x_F)p_T + C(x_F)p_T^2) e^{-D(x_F)p_T^{E(x_F)}}$$

$$e.g. B = \sum_i b_i x_F^i \quad x_F = \frac{p_Z}{120}$$

- New N24LE parametrization
- BCDE parameters are polynomials in x_F
- Empirical approach in different regions of p_T
- BCDE are then parameters in fit to data

Focusing Parameter





- LE (top) and ME (bottom) energy peak mismatch between MC and data
- This was considered to be due to the simulation of the focusing
- Horn Current Miscalibration parameter was used to correct peak
- Preferred horn current correction is so large (~10kA) that it is labelled an **effective parameter** incorporating all beam optics

The Horn Off sample

- Runs with no current through the magnetic focusing horns
- Low stats, but virtually unaffected by beam systematics
- They can be used to constrain hadron production parameters
- Useful to fit with horn on runs, as all samples contain 'horn-off-type' events

Region of high p_z low p_T 'horn off type' events, largely unaffected by magnetic focussing



Effective parameters for ND

- There is also unknown physics in the ND
- Normalization based on neutrino interaction is introduced to pick up ND effects

QE

$$\nu_{\mu} + n \rightarrow \mu^{-} + p$$


Resonance

$$\nu_{\mu} + p \rightarrow \mu^{-} + \Delta^{++} \rightarrow \mu^{-} + p + \pi^{+}$$

DIS

$$\nu_{\mu} + N \rightarrow \mu^{-} + \textit{hadronic shower}$$

R1: Data, and MC split into interaction components



Extracting information from the fits





Target Parameters	Focusing Parameters	ND Parameters
<ul style="list-style-type: none">• Hadron Production• Target Decay• Target Position• Intensity	<ul style="list-style-type: none">• Focusing Horn Current• Decay pipe atmosphere	<ul style="list-style-type: none">• Interaction Effects

Fit Results









- Fit with different target position samples + horn off to cover wide range of p_T p_Z space and constrain focusing systematics, obtaining ND parameters



MINOS preliminary

Applying parameters to other samples


- Parameters found in previous fit used to reweight different MC, results in moderately improved fit to data



R2&3 target decay accounted for

Testing the ND physics model


- Random Phase Approximation (RPA) weights are used in the simulation
- This models the effect of complex inter-nucleon interactions in atoms on the neutrino cross section in the Near Detector
- The overall effect of the change in potential is a suppression of quasi-elastic type neutrino interactions at low energy [3]



Parameter reaction to RPA weights



Fractional change in Parameters, RPA and no RPA



- We are trying to improve understanding the components of the NuMI beam simulation, due to its relevance in contemporary neutrino beams
- Improved hadron production momentum parametrisation
- Use effective parameters to separate Target, Focussing, and detector interaction effects during fit to data
- Parameter results shown to improve the MC of other MINOS run configurations
- Can hopefully be used to inform beam design + simulation

Invite Questions




UCL



Backup

Beam systematics

- Also incorporate effects associated with other beam components
- Some are fixed based on surveys – Target Z position
- Others are free parameters in the fit – Horn Current Miscalibration, target decay



Schematic of the NuMI beam, reproduced from [2]


Parameters RPA v no RPA




	RPA	No RPA
Pi 0 :	0.649422 +/- 0.092940	0.663331 +/- 0.086148
Pi 1 :	3.237669 +/- 0.303910	3.258092 +/- 0.213383
Pi 2 :	1.163347 +/- 0.028383	1.164678 +/- 0.013039
Pi 3 :	-1.335198 +/- 0.279268	-1.312645 +/- 0.088701
I mis :	-6.062433 +/- 0.215879	-6.017456 +/- 0.205155
QE norm :	1.401484 +/- 0.208088	1.065699 +/- 0.170377
RES norm :	1.327494 +/- 0.168526	1.383580 +/- 0.139345
DIS norm :	1.038960 +/- 0.035984	1.042975 +/- 0.016188
other norm :	1.000000 +/- 0.000000	1.000000 +/- 0.000000


Parameters RPA v no RPA

RPA v no RPA actual parameters






Pt Pz binning scheme




Typical Fit to data

- Results of a joint MC fit to the data with Run 1 and horn off samples
- 8 hadron production parameters + 1 focusing parameter
- Minuit is used to fit MC to data using parameters


Run 1 Data and MC




R1 Data/MC ratio



Horn off




Horn off data/mc ratio





Effect of RPA weights on Hadron Production Parameters (R1 & Horn Off fit)



Fit Results

- Normalisations of interaction types do not vary hugely as hadron production parameters are introduced, altering target weightings

