

Status and latest results of the ν GeN experiment at the Kalinin Nuclear Power Plant


Ignatov Georgii for the ν GeN collaboration

The study was supported by the Russian Science Foundation (project no. 24-72-10089)

LPI RAS, MIPT

22.07.2025

21st Rencontres du Vietnam, Neutrino physics



- **Coherent Elastic ν - Nucleus Scattering:** was predicted in 1974^{1,2} and observed for the first time in 2017³
- SM cross-section:

$$\frac{d\sigma}{dT} = \frac{G_F^2 M}{4\pi} ([1 - 4 \sin^2 \theta_W]Z - N)^2 \left(1 - \frac{T}{T_{max}}\right) F^2(q^2)$$

$$T_{max} = \frac{2E_\nu^2}{M + 2E_\nu}$$

Nucleus	T_{max} , keV ($E_\nu = 5$ MeV)	T_{max} , keV ($E_\nu = 30$ MeV)
^{12}C	4.44	159.0
^{23}Na	2.32	83.2
^{40}Ar	1.33	47.9
^{74}Ge	0.72	25.9
^{133}Cs	0.40	14.4

Motivation:


- NC ν -q NSI / SM test
- Nuclear FF
- BG for DM search
- Supernova dynamics
- Reactor monitoring

¹ D. Freedman, PRD v.9, iss.5 (1974)

² V.Kopeliovich, L. Frankfurt, ZhETF. Pis. Red., v.19 n.4 (1974)

³ Akimov, D., et al. Science 357.6356 (2017): 1123-1126.

CE ν NS search




Ge CE ν NS tension

- Dresden-II (2022): observation claim for D1/D2 QF
- COHERENT (2024/2025): 2σ below SM prediction
- CONUS+ (2025): observation claim for C QF

ν GeN experiment

- KNPP unit 3 (~ 50 m.w.e. overburden).
- ≈ 11 distance from the center of the reactor core.
- antineutrino flux $- 4.4 \times 10^{13} \text{ cm}^{-2} \text{ s}^{-1}$.
- Reactor cycle: 16.5 months ON, 45 days OFF.




Main goals of the ν GeN experiment:

- CE ν NS measurement
- neutrino EM properties search

ν GeN setup




- 1.4 kg active mass HPGe PPC.
- 102 eV FWHM from pulser.
- 0.29 keV threshold.
- Cooled by cryocooler.



- Multi-layered shielding + muon veto.
- Anti-vibration platform.
- Comparison of amplifiers with different shaping times allows noise discrimination.
- Total exposition: $\sim 2000 \text{ kg} \times \text{d}$ up to 2025, but different conditions.

Dataset & stability


- Data collected from October 2022 to May 2023.
- 38 d OFF (black) & 137 d ON (red).
- Stability checked by the intensity of Rn/neutron-induced lines, cosmogenic lines and total count rate in ROI.



Quenching factor (QF)

$$QF(T_{nr}) = \frac{E_{det}(T_{nr})}{E_{det}(T_{er})} \Big|_{T_{nr}=T_{er}}$$




Discrepancies in measurements of QF for low energies:



3 scenarios: CONUS (C), Dresden FeF-based (D1) and photo-neutron based (D2).

Fit & results

QF scenarios: C – red, D1 – magenta, D2 – green.




QF	$A_{best} \pm \sigma_A, \times \text{SM}$	χ_{best}^2 (ndf=10)	S, $\times \text{SM}$	L, $\times \text{SM}$
C	1.5 ± 1.7	13.6	3.8	4.3
D1	0.1 ± 0.4	14.4	1.6	0.7
D2	0.8 ± 1.4	14.1	3.3	3.1

Sensitivity: Full statistics + BG model \rightarrow
null-hypothesis rejection at $\sim 5\sigma$.

- Upper limit (90% C.L.):
 $4.3 \times \text{SM}$ (C QF)
- Tension with D1 QF

ν GeN modifications

”Compton veto”: ~ 30 kg NaI




More than 2 times BG suppression in the lab with shielding & μ veto. Now tests at the Baksan underground laboratory.



- DAQ modification: waveform analysis \rightarrow noise / surface events rejection.
- Cryocooler modification: reduce power consumption \rightarrow less vibrations \rightarrow less noise.

ν EM properties




ν - γ vertex:

$$\Lambda_\mu(q) = F_Q(q^2)\gamma_\mu - F_M(q^2)i\sigma_{\mu\nu}q_\nu + F_E(q^2)\sigma_{\mu\nu}q_\nu\gamma_5 + F_A(q^2)(q^2\gamma_\mu - q_\mu q_\nu\gamma_\nu)\gamma_5$$


Observables: Magnetic Moment (MM), (milli)charge, charge radii.

NMM in SM extensions*:




*SJ, PoS(DISCRETE2020-2021)037

ν -electron CS for $E_\nu = 5$ MeV:




Limits on ν EM properties


All limits are given for 90% C.L.; ROI: [0.4,17] keV & +[20,60] keV for NMM.



$$\mu_\nu < 7.5 \times 10^{-11} \mu_B$$



$$q_{ij} < 2.4 \times 10^{-12} e$$



$$q_{ii} \in (-2.2, 2.3) \times 10^{-12} e$$

Limit, $10^{-11} \mu_B$	Experiment
7.5	CONUS
7.5	ν GeN
7.4	TEXONO
2.9	GEMMA
0.64	XENONnT

Limit, $10^{-12} e$	Experiment
2.7	GEMMA
1.2	TEXONO
2.4 (0.9)	ν GeN
0.6	CONUS+
0.224	LZ

Astrophys.: $\mu_\nu < 1.2 \times 10^{-12} \mu_B$

Cosmology: $q_\nu \leq 10^{-35} e$

Summary

CEvNS search [V. Belov et al. Chinese Phys. C 49 053004 (2025)]:

- Limit on the CEvNS count rate at $4.3 \times \text{SM}$ (C QF).
- Tension with Dresden-II QF and CEvNS claim.

Limits on ν EM properties:

- $\mu_\nu < 7.5 \times 10^{-11} \mu_B$.
- $q_\nu < 2.4 (0.9) \times 10^{-12} e$ for FEA (EPA) CS.

+ Ongoing analysis to use full statistics ($\sim 2000 \text{ kg} \times \text{d}$) and upgrades to reduce BG + lower threshold.




Thank you for your attention!




Backup

Stability: neutrons, Rn


Neutrons in BC501A LS outside shielding: First estimation: $\sim 10^{-2}$ of BG rate.



Rn by ^{214}Bi 609 keV line:



Sensitivity estimation



«Time» corresponds to ON / OFF statistics depending on the analysis strategy.