

# Physics with Reactor Neutrinos

Junting Huang

Shanghai Jiao Tong University

July 20, 2023

Vietnam School on Neutrinos


ICISE center, Quy Nhon, VN



# Table of Contents

1. Introduction to Reactor Neutrinos
2. Discovery of Neutrinos: Cowan and Reines Experiment
3.  $\theta_{12}$  and  $\Delta m_{21}^2$  Measurement: KamLAND Experiment
4.  $\theta_{13}$  Measurement: Daya Bay, RENO and Double CHOOZ
5. Sterile Neutrino Searches: STEREO, etc.
6. Mass Ordering Measurement: JUNO Experiment
7. Summary

# Research Experience



# Neutrino Sources





Figure 1: Neutrino sources and their energies. Figure taken from [1].


# Nuclear Reactor

- ▶ nuclear fission, first reactor is Chicago Pile-1, Fermi, 1942
- ▶ a typical nuclear reactor produces  $\mathcal{O}(1 \text{ GW})$  of electricity




# Reactor Neutrinos

- ▶ 99.9% of  $\bar{\nu}_e$  from fission isotopes of  $^{235}\text{U}$ ,  $^{238}\text{U}$ ,  $^{239}\text{Pu}$ ,  $^{241}\text{Pu}$
- ▶ per fission decay chain: 200 MeV, 6 neutrinos, pure  $\bar{\nu}_e$
- ▶  $2 \times 10^{20} \text{ GW}^{-1} \text{ s}^{-1}$  reactor neutrinos



# Neutrino Flux and Neutrino Detection



- ▶ inverse-beta decay (IBD): prompt signal,  $E_{e^+} = E_\nu - 0.8 \text{ MeV}$
- ▶ neutron capture: delayed signal

# Fuel Evolution

- ▶ evolution of reactor core fuel
- ▶ antineutrino flux depends on fission fraction





Figure 2: Evolution of reactor core fuel in Daya Bay [2].



# Discovery of Neutrinos: Cowan and Reines Experiment

# Original Plan of Cowan and Reines to Detect Neutrinos




**Figure 1. Detecting Neutrinos from a Nuclear Explosion**


Antineutrinos from the fireball of a nuclear device would impinge on a liquid scintillation detector suspended in the hole dug below ground at a distance of about 40 meters from the 30-meter-high tower. In the original scheme of Reines and Cowan, the antineutrinos would induce inverse beta decay, and the detector would record the positrons produced in that process. This figure was redrawn courtesy of Smithsonian

## Early Reactor Neutrino Measurement

- ▶ neutrino rate: from  $10^{18} \text{ cm}^{-2} \text{ s}^{-1}$  near atomic bomb site, to  $10^{14} \text{ cm}^{-2} \text{ s}^{-1}$  at a reactor
- ▶ initial measurement using the nuclear reactors at Hanford (Washington State) in 1953
- ▶ inverse beta decay




- ▶ liquid scintillator (LS) with cadmium dissolved in it



- ▶ coincidence of positron annihilation and neutron capture


## Detector at Hanford

- ▶ 28 inches in diameter, 30 inches in height, 300-liter capacity
- ▶ 90 2-inch PMTs, “Herr Auge”



**Figure 3:** The scintillation detector for the 1953 neutrino detection experiment at Hanford.

# Detecting IBD





## Moving to Savannah River

- ▶ “it was felt that an identification of the free neutrino had probably been made.”
- ▶ moved experiment to Savannah River Plant (South Carolina)
- ▶ a detector tank containing more than 1 ton of scintillation fluid in a crawling convoy of 5 over-sized trucks




# Experimental Setup at Savannah River



- ▶ club-sandwich setup
- ▶ bread layers: 0.6 m thick, 1.9 m long, 1.4 m wide
  - ▶ triethylbenzene solution of terphenyl (scintillator  $C_{18}H_{14}$ ) and POPOP (wavelength shifter)
  - ▶ 110 5-inch PMTs in total, res. at 0.5 MeV is 30% FWHM
- ▶ meat layers: water solution of cadmium chloride
- ▶ parafin and lead shield, underground room

## Data Acquisition


- ▶ trigger on pulse amplitudes and coincidences
- ▶ pulses on the triple-beam oscilloscopes were recorded photographically
- ▶ short delay (up to  $17\ \mu\text{s}$ ): good signal-to-background ratio
- ▶ calibrations: copper-64 positron source, plutonium-beryllium neutron source, and cosmic ray






## Background

- ▶ pulses in all three traces





- ▶ strange signals



- ▶ accidental background: radio-activities from materials

# Cowan and Reines




## Experimental Results

- ▶ total running time: 1371 hours = 57 days
- ▶ cross section  $6.3 \times 10^{-44} \text{ cm}^2$  (about right)
- ▶ in one run, neutrino signal rate was 0.6 counts per hour, signal is 20 times the accidental background due to reactor



## 1995 Nobel Prize


- ▶ Cowan passed way at 54 in 1974. Reines received Nobel price in 1995 in both their names.
- ▶ “for pioneering experimental contributions to lepton physics” jointly with one half to Martin L. Perl “for the discovery of the tau lepton” and with one half to Frederick Reines “for the detection of the neutrino”
- ▶ among 79 / 116 Nobel Prizes in particle physics (1901—2022)




$\theta_{12}$  and  $\Delta m_{21}^2$  Measurement:  
KamLAND Experiment

# Discovery of Neutrino Oscillations

- ▶ Super-Kamiokande: atmospheric  $\nu_\mu$  disappearance
- ▶ SNO:  $\nu_e$  flux and total neutrino  $\nu_{e,\mu,\tau}$  flux from the Sun
- ▶ neutrinos can change flavor as they travel
- ▶ non-zero masses, 2015 Nobel Prize



(a) Super-Kamiokande, 1998 [3]




(b) SNO, 2002 [4]

# Standard Neutrino Oscillation Model

- ▶  $\nu_\alpha$ : state with specific flavor ( $\nu_e, \nu_\mu, \nu_\tau$ )
- ▶  $\nu_i$ : state with specific mass ( $m_1, m_2, m_3$ )
- ▶  $|\nu_\alpha\rangle = \sum_i U_{\alpha i} |\nu_i\rangle$ , matrix  $U$  is unitary, called PMNS matrix

$$U = \begin{pmatrix} 1 & & \\ & c_{23} & s_{23} \\ & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} & c_{13} & -s_{13}e^{i\delta_{CP}} \\ & 1 & \\ -s_{13}e^{i\delta_{CP}} & & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} \\ -s_{12} & c_{12} \\ & & 1 \end{pmatrix}$$


- ▶ 2-flavor example:  $P(\nu_\alpha \rightarrow \nu_\beta) = \sin^2(2\theta) \sin^2 \left[ \Delta m_{ij}^2 (L/4E) \right]$   
 $\Delta m_{ij}^2 = m_i^2 - m_j^2$ ,  $L$  is baseline,  $E$  is energy



# Reactor Neutrino Oscillations

- ▶ for reactor neutrinos:  $\bar{\nu}_e \rightarrow \bar{\nu}_\mu$  or  $\bar{\nu}_\tau$
- ▶ oscillations depend on  $\theta_{12}$ ,  $\theta_{13}$ ,  $\Delta m_{21}^2$ ,  $\Delta m_{31}^2$


$$P(\nu_e \rightarrow \nu_e) \approx 1 - \sin^2(2\theta_{12}) \sin^2\left(\frac{\Delta m_{21}^2 L}{4E}\right) - \sin^2(2\theta_{13}) \sin^2\left(\frac{\Delta m_{31}^2 L}{4E}\right)$$






# KamLAND Experiment

- ▶ KamLAND: Kamioka Liquid Scintillator Anti-Neutrino Detector, proposed in 1994, started taking data in 2002
- ▶ average baseline of 180 km



- ▶ first oscillation observation from man-made neutrino
- ▶  $\Delta m_{21}^2$  was determined to a high precision


# KamLAND Detector




- ▶ about 2000 PMTs, 34% photocathode coverage
- ▶ 18 m diameter vessel, 13 m diameter, 1 kton LS
- ▶ energy resolution:  $6.4\%/\sqrt{E(\text{MeV})}$ , position resolution  $12\text{ cm}/\sqrt{E(\text{MeV})}$

## Backgrounds in KamLAND Experiment

- ▶ unexpected:  $^{13}\text{C}(\alpha, n)^{16}\text{O}$ , liquid scintillator purification
- ▶  $\text{geo-}\bar{\nu}_e$ : cut on 2.6 MeV prompt signal




- ▶  $^9\text{Li}$  and  $^8\text{He}$ : 2 s veto of entire volume for showering muon or a 2 s veto of a 3-m-radius volume around a muon track.







## KamLAND Data in $L/E$

- ▶ ratio of the observed  $\bar{\nu}_e$  spectrum to the expectation for no-oscillation as a function of  $L/E$  [5]
- ▶ three flavor fit, strong constraint on  $\Delta m_{21}^2$





# Oscillation Results from KamLAND



# KamLAND-Zen

- ▶ in 2011,  $> 300 \text{ kg } ^{136}\text{Xe}$  loaded in 13 ton liquid scintillator
- ▶ KamLAND-800: 745 kg Xe, 91% enriched, loaded with 3.1%  $^{110m}\text{Ag}$  from Fukushima (2011)
- ▶ 270 keV FWHM




$\theta_{13}$  Measurement: Daya Bay, RENO and  
Double CHOOZ

# Significance of $\theta_{13}$ Measurement

- ▶ non-zero value of  $\theta_{13}$  allows the measurement of  $\delta_{CP}$ , matter-antimatter asymmetry of the universe

$$U = \begin{pmatrix} 1 & & \\ & c_{23} & s_{23} \\ & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & & -s_{13}e^{i\delta_{CP}} \\ & 1 & \\ -s_{13}e^{i\delta_{CP}} & & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} \\ -s_{12} & c_{12} \\ & & 1 \end{pmatrix}$$

- ▶ indications of a nonzero  $\theta_{13}$  in 2011: T2K [6], MINOS [7], Double Chooz [8]




- ▶ excluded non-zero  $\theta_{13}$  above  $5\sigma$  in 2012: Daya Bay and RENO




# Near and Far Detectors of Daya Bay Experiment


- ▶ 2 nuclear power plants (NPP), 6 reactor cores
- ▶ 4 far detectors (EH3) and 4 near detectors (EH1 and EH2)




# Detector Design



# Daya Bay Detectors in EH3



## Interior of a Daya Bay Detector



# Daya Bay Results in 2012 [9]

- ▶ exposure of 55 days, rate measurement
- ▶ determine non-zero  $\theta_{13}$  above 5 sigma level




Figure 5: Daya Bay experimental results in 2012 [9].

## Daya Bay Results in 2023 [10]

- ▶  $\sin^2 2\theta_{13} = 0.0851 \pm 0.0024$ ,
- ▶ “The reported  $\sin^2 2\theta_{13}$  will likely remain the most precise measurement of  $\theta_{13}$  in the foreseeable future”
- ▶ “crucial to the investigation of the mass hierarchy and CP violation in neutrino oscillation”




Figure 6: Daya Bay experimental results in 2023 [10].

# RENO Experiment

- ▶ two detectors: near (290 m) and far detectors (1380 m)
- ▶ 16.5 t, 0.1% Gd-doped liquid scintillator
- ▶ pure LS ( $\gamma$  catcher), mineral oil (shielding), water
- ▶ 229 days exposure, first results in 2012 at  $> 4\sigma$





Figure 7: Detector locations and design in RENO experiment [11].

# Double CHOOZ Experiment

- ▶ single detector operation: April 2011 until January 2013
- ▶ both detectors FD and ND: January 2015 until April 2016




Figure 8: Baseline and detectors of Double CHOOZ experiment [12].



# A Comparison of $\theta_{13}$ Measurements





Figure 9: A comparison on  $\theta_{13}$  measurements in 2020 [12]. Double CHOOZ 2021 [13] and Daya Bay 2023 results [10] not included.

Sterile Neutrino Searches: STEREO, etc.


# Reactor Antineutrino Anomaly

- ▶ predicting  $\bar{\nu}_e$  flux from beta spectra for  $^{235}\text{U}$ ,  $^{239}\text{Th}$  and  $^{241}\text{Th}$
- ▶ Huber–Mueller (HM) model in 2011, measured rates about 5% below prediction, **reactor antineutrino anomaly**
- ▶ sterile neutrinos:  $P = 1 - \sin^2(2\theta) \sin^2(\Delta m_{41}^2 L/4E)$




## 5 MeV “Bump”

- ▶ a “bump” at 5–6 MeV, with an amplitude of about 10% [12]
- ▶ seen in the data of Daya Bay, RENO, Double CHOOZ, and NEOS experiments





# STEREO Experiment [14]

- ▶ highly enriched  $^{235}\text{U}$  fuel, 10 m baseline, 58 MW research reactor at ILL, France
- ▶ shielding: lead, borated polystyrene, mu-metal
- ▶ started taking data in 2016, 400 neutrino events per day




# STEREO Results

- ▶ Nature 613, 257–261 (2023), excluded most of the reactor anomaly parameter space
- ▶ agree with the prediction of Letourneau et al. [15]



# Calculations by Letourneau et al. in 2022

- ▶ Phys. Rev. Lett. 130, 021801 (2023) [15], summation model
- ▶ with a single parameter, can reproduce both norm and shape
- ▶ suspect that the anomalous feature could be from a shape bias in the  $\beta^-$  energy spectra measured at ILL



## Other Very Short Baseline Experiments

- ▶ NEOS: Yeonggwang, Korea, 24 m [16]
- ▶ PROSPECT: Oak Ridge,  $^{235}\text{U}$  fissions, 8 m,  $^6\text{Li}$ -doped LS [17]
- ▶ Neutrino-4, DANSS, etc. [18]





Figure 10: NEOS (left plot, 2017, [16]) and PROSPECT (right plot, 2021, [17]) results on sterile neutrinos.



# Mass Ordering Measurement: JUNO Experiment

# JUNO Experiment

- ▶ JUNO: Jiangmen Underground Neutrino Observatory
- ▶ reactor  $\bar{\nu}_e \rightarrow \bar{\nu}_e$ : neutrino mass ordering



- ▶ sub-percent measurement of  $\theta_{12}$ ,  $\Delta m_{21}^2$  and  $|\Delta m_{31}^2|$
- ▶ multi-purpose detector: geoneutrinos, atmospheric neutrinos, solar neutrinos, supernova neutrinos

# Detector Location and Baseline

- ▶ located at southern China, near Kaiping, Guangdong province
- ▶ two nuclear power plants (NPP), 53 km baseline



Summary of the thermal power and baseline to the JUNO detector for the Yangjiang (YJ) and Taishan (TS) reactor cores, as well as the remote reactors of Daya Bay (DYB) and Huizhou (HZ).

Cores	YJ-1	YJ-2	YJ-3	YJ-4	YJ-5	YJ-6	TS-1	TS-2	DYB	HZ
Power (GW)	2.9	2.9	2.9	2.9	2.9	2.9	4.6	4.6	17.4	17.4
Baseline(km)	52.74	52.82	52.41	52.49	52.11	52.19	52.77	52.64	215	265


# JUNO Experimental Site

- ▶ vertical tunnel and slope tunnel (42°)
- ▶ overburdens 650 m (1800 w.m.e.), about 4 Hz muons rate




# JUNO Detector

- ▶ 20 kton liquid scintillator: LAB + PPO + bis-MSB
- ▶ top tracker and water pool for muon veto



# JUNO PMTs

- ▶ 3% energy resolution at 1 MeV, 78% photocathode coverage
- ▶ 17612 20 in PMTs: NNVT and Hamamatsu
- ▶ 25600 3 in PMTs: for large PMT non-linearity correction




17612 (CD) + 2400 (Veto) 20-inch PMTs



25600 3-inch PMTs

# Detector Frame



# Acrylic Sphere






# PMT Array



By [Jagjit](#) on 2025-04-23 13:52


# Energy Spectra

- ▶ signal: about 42 events per day
- ▶ major background: accidentals, geoneutrinos,  ${}^9\text{Li}/{}^8\text{He}$ , global reactors, with about 1 event per day each




# Mass Ordering Sensitivity

- ▶ data taking will start in 2024
- ▶ TAO, satellite detector by Taishan reactors, flux constraint
- ▶  $3\sigma$  (reactors only) with about 6 years  $\times$  26.6 GW exposure



# Precision Measurement of Oscillation Parameters

- ▶ precision of  $\sin^2 \theta_{12}$ ,  $\Delta m_{21}^2$  and  $|\Delta m_{31}^2| < 0.5\%$  in 6 years [19]
- ▶ current uncertainty from PDG 2020: 4.2% for  $\sin^2 \theta_{12}$ , 2.4% for  $\Delta m_{21}^2$  and 1.3% for  $|\Delta m_{31}^2|$



# Summary

- ▶ reactor neutrinos:  $\bar{\nu}_e$  from  $^{235}\text{U}$ ,  $^{238}\text{U}$ ,  $^{239}\text{Pu}$ , and  $^{241}\text{Pu}$ , inverse beta decay, Cd/Gd/Li for neutron capture
- ▶ **discovery of neutrinos**: Cowan-Reines experiment, 1956
- ▶  $\theta_{12}$  and  $\Delta m_{21}^2$ : KamLAND experiment 2002–2011
- ▶  $\theta_{13}$ : Daya Bay, RENO, Double CHOOZ
- ▶ **sterile neutrinos**: STEREO, etc., normalization and shape anomaly may have been solved
- ▶ **mass ordering**: JUNO,  $3\sigma$  after 6 years starting 2024
- ▶ other topics not discussed: nuclear security, etc.

## References I

- <sup>1</sup> J. L. Hewett et al., “Planning the Future of U.S. Particle Physics (Snowmass 2013): Chapter 2: Intensity Frontier”, in *Community Summer Study 2013: Snowmass on the Mississippi* (Jan. 2014).
- <sup>2</sup> F. P. An et al., “Evolution of the Reactor Antineutrino Flux and Spectrum at Daya Bay”, *Phys. Rev. Lett.* **118**, 251801 (2017).
- <sup>3</sup> Y. Fukuda et al., “Evidence for oscillation of atmospheric neutrinos”, *Phys.Rev.Lett.* **81**, 1562–1567 (1998).
- <sup>4</sup> Q. Ahmad et al., “Direct evidence for neutrino flavor transformation from neutral current interactions in the Sudbury Neutrino Observatory”, *Phys.Rev.Lett.* **89**, 011301 (2002).
- <sup>5</sup> M. P. Decowski, “KamLAND’s precision neutrino oscillation measurements”, *Nucl. Phys. B* **908**, 52–61 (2016).

## References II

- <sup>6</sup> K. Abe et al., “Indication of Electron Neutrino Appearance from an Accelerator-produced Off-axis Muon Neutrino Beam”, *Phys. Rev. Lett.* **107**, 041801 (2011).
- <sup>7</sup> P. Adamson et al., “Improved search for muon-neutrino to electron-neutrino oscillations in MINOS”, *Phys. Rev. Lett.* **107**, 181802 (2011).
- <sup>8</sup> Y. Abe et al., “Indication for the disappearance of reactor electron antineutrinos in the Double Chooz experiment”, *Phys.Rev.Lett.* **108**, 131801 (2012).
- <sup>9</sup> F. An et al., “Observation of electron-antineutrino disappearance at Daya Bay”, *Phys.Rev.Lett.* **108**, 171803 (2012).
- <sup>10</sup> F. P. An et al., “Precision Measurement of Reactor Antineutrino Oscillation at Kilometer-Scale Baselines by Daya Bay”, *Phys. Rev. Lett.* **130**, 161802 (2023).

## References III

- <sup>11</sup>J. Ahn et al., “Observation of Reactor Electron Antineutrino Disappearance in the RENO Experiment”, *Phys.Rev.Lett.* **108**, 191802 (2012).
- <sup>12</sup>H. de Kerret et al., “Double Chooz  $\theta_{13}$  measurement via total neutron capture detection”, *Nature Phys.* **16**, 558–564 (2020).
- <sup>13</sup>T. Abrahão et al., “Reactor rate modulation oscillation analysis with two detectors in Double Chooz”, *JHEP* **01**, 190 (2021).
- <sup>14</sup>H. Almazán et al., “STEREO neutrino spectrum of  $^{235}\text{U}$  fission rejects sterile neutrino hypothesis”, *Nature* **613**, 257–261 (2023).
- <sup>15</sup>A. Letourneau, V. Savu, D. Lhuillier, T. Lasserre, T. Materna, G. Mention, X. Mougeot, A. Onillon, L. Perisse, and M. Vivier, “Origin of the Reactor Antineutrino Anomalies in Light of a New Summation Model with Parametrized  $\beta$ - Transitions”, *Phys. Rev. Lett.* **130**, 021801 (2023).



## References IV

- <sup>16</sup>Y. J. Ko et al., “Sterile Neutrino Search at the NEOS Experiment”, *Phys. Rev. Lett.* **118**, 121802 (2017).
- <sup>17</sup>M. Andriamirado et al., “Improved short-baseline neutrino oscillation search and energy spectrum measurement with the PROSPECT experiment at HFIR”, *Phys. Rev. D* **103**, 032001 (2021).
- <sup>18</sup>C. Awe et al., “High Energy Physics Opportunities Using Reactor Antineutrinos”, (2022).
- <sup>19</sup>A. Abusleme et al., “Sub-percent precision measurement of neutrino oscillation parameters with JUNO”, *Chin. Phys. C* **46**, 123001 (2022).

## $\bar{\nu}_e$ Disappearance

$$\begin{aligned}
 P(\nu_e \rightarrow \nu_e) &= 1 - \sin^2(2\theta_{12}) \cos^4 \theta_{13} \sin^2\left(\frac{\Delta_{21}}{2}\right) \\
 &\quad - \sin^2(2\theta_{13}) \left[ \cos^2 \theta_{12} \sin^2\left(\frac{\Delta_{31}}{2}\right) + \sin^2 \theta_{12} \sin^2\left(\frac{\Delta_{32}}{2}\right) \right] \\
 &\approx 1 - \sin^2(2\theta_{12}) \sin^2\left(\frac{\Delta_{21}}{2}\right) - \sin^2(2\theta_{13}) \sin^2\left(\frac{\Delta_{31}}{2}\right)
 \end{aligned}$$

