

VIETNAM SCHOOL ON NEUTRINOS (VSON 7)

# Introduction to Neutrino Interactions

**Nguyen Thi Hong Van**

Institute of Physics

Vietnam Academy of Science and Technology

# References

- K Zuber, “*Neutrino Physics*”, Institute of Physics Publishing, 2004.
- C. Giunti and C.W.Kim, “*Fundamentals of Neutrino Physics and Astrophysics*”, Oxford University Press, 2007.
- Ulrich Mosel, “*Neutrino Interactions with Nucleons and Nuclei: Importance for Long-Baseline Experiments*”, arXiv:1602.00696v3
- Kevin McFarland, “*Neutrino Interactions*”, arXiv:0804.3899v1
- Kevin McFarland, “Interactions of Neutrinos”, lectures at INWS 2015, Brasil, August 2015.
- Steve Boyd, Lecture of “*Neutrino Physics*”
- Neutrino Interactions - Thesis - Daniel I. Scully

# Contents

- Introduction to neutrinos and neutrino interactions
- Weak interaction in Standard Model
- Weak interaction for neutrinos
- Neutrino – Electron scattering
- Neutrino – Nucleons scattering
- Neutrino – Nucleus scattering
- Neutrino Event Generators

# Introduction to neutrinos

## Neutrinos

No electric charge

No color

Participate in weak interaction

Cannot be observed  
Directly in detector

## Standard Model of Elementary Particles

	three generations of matter (fermions)			interactions / force carriers (bosons)	
	I	II	III		
mass	$\approx 2.2 \text{ MeV}/c^2$	$\approx 1.28 \text{ GeV}/c^2$	$\approx 173.1 \text{ GeV}/c^2$	0	$\approx 124.97 \text{ GeV}/c^2$
charge	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	0	0
spin	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	0
	<b>u</b> up	<b>c</b> charm	<b>t</b> top	<b>g</b> gluon	<b>H</b> higgs
	<b>d</b> down	<b>s</b> strange	<b>b</b> bottom	<b><math>\gamma</math></b> photon	
	<b>e</b> electron	<b><math>\mu</math></b> muon	<b><math>\tau</math></b> tau	<b>Z</b> Z boson	
	<b><math>\nu_e</math></b> electron neutrino	<b><math>\nu_\mu</math></b> muon neutrino	<b><math>\nu_\tau</math></b> tau neutrino	<b>W</b> W boson	

**QUARKS** (left side of the table)

**LEPTONS** (left side of the table)

**GAUGE BOSONS VECTOR BOSONS** (bottom right)

**SCALAR BOSONS** (right side)

# Why study neutrino interactions?

- Neutrino interaction is a signature of neutrino existence
- Study neutrino interactions to understand the weak interaction and electroweak unification theories.
- Neutrino interactions give information for neutrino mass and neutrino mixing,...
- In experiments: better understanding of neutrino interactions → more precise measuring neutrino oscillations.

# Weak Interactions of SM

$$L_{\text{int}} = i \frac{g}{\sqrt{2}} [j_{\mu}^{(+)} W^{\mu} + j_{\mu}^{(-)} W^{\mu+}] + i \frac{g}{2 \cos \theta_W} j_{\mu}^{(Z)} Z^{\mu}$$

- Charged Current (CC) interactions:  
(mediated by  $W$  bosons)
- Neutral Current (NC) interaction:  
(mediated by  $Z$  boson)

- Weak mixing angle:

$$\frac{g_W}{g_Z} = \cos \theta_W$$

Coupling strength

$$j_{\mu}^{\pm} = \bar{u} \frac{-ig_W}{2\sqrt{2}} \gamma^{\mu} (1 \pm \gamma^5) u$$


Dirac spinors

$$j_{\mu}^0 = \bar{u} \frac{-ig_Z}{2} (g_V \gamma^{\mu} - g_A \gamma^{\mu} \gamma^5) u$$

Particles	$g_V$	$g_A$
Neutrinos	$\frac{1}{2}$	$\frac{1}{2}$
Charged Leptons	$\frac{1}{2} + 2 \sin^2 \theta_W$	$-\frac{1}{2}$
Up-type Quarks	$\frac{1}{2} - \frac{4}{3} \sin^2 \theta_W$	$\frac{1}{2}$
Down-type Quarks	$-\frac{1}{2} + \frac{2}{3} \sin^2 \theta_W$	$-\frac{1}{2}$

# Parity violation by weak interaction


# Parity symmetry



$$\begin{aligned}\vec{E}(\vec{x}, t) &\xrightarrow{P} -\vec{E}(-\vec{x}, t) \\ \vec{B}(\vec{x}, t) &\xrightarrow{P} \vec{B}(-\vec{x}, t) \\ \vec{j}(\vec{x}, t) &\xrightarrow{P} -\vec{j}(-\vec{x}, t) \\ \nabla &\xrightarrow{P} -\nabla\end{aligned}$$



# Example of Parity symmetry violation



World  $\neq$  Mirror World  
(parity violation)

Lecture for CERN summer student  
Given by Tatsuya Nakada 9

# Parity violation by weak interaction

- Vector current (*odd parity*):  $\hat{P}(\bar{\psi}\gamma^\mu\psi) = -(\bar{\psi}\gamma^\mu\psi)$
- Axial-vector current (*even parity*):  $\hat{P}(\bar{\psi}\gamma^\mu\gamma^5\psi) = \bar{\psi}\gamma^\mu\gamma^5\psi$
- Weak current is a mixture of vector and axial vector currents:

$$j_\mu^\pm = \bar{u} \frac{-ig_W}{2\sqrt{2}} \gamma^\mu (1 \pm \gamma^5) u$$



- → Parity of a system is violated by weak force! (first postulated by Lee & Yang in 1950)

# Parity violation confirmed by experiment

- **Helicity operator:** projection of spin on the momentum direction.

$$H = \frac{\vec{\sigma} \cdot \vec{p}}{|\vec{p}|} \xrightarrow{P} \frac{\vec{\sigma} \cdot (-\vec{p})}{|\vec{p}|} = -H$$

- Parity violation appeared in the the asymmetry of polarized  $^{60}\text{Co}$  (by Wu in 1957):



More electrons emitted  
in direction opposite to  
 $^{60}\text{Co}$  spin  $\rightarrow$  parity violation!

# Weak interaction and chirality

- CC weak interaction:

$$j_{\mu}^{\pm} = \bar{u} \frac{-ig_W}{2\sqrt{2}} \gamma^{\mu} (1 \pm \gamma^5) u$$

- Chirality:

Lorentz invariant

but not directly measurable!:

$$u = u_L + u_R$$

$$u_L = \frac{1}{2} (1 - \gamma^5) u$$

$$u_R = \frac{1}{2} (1 + \gamma^5) u$$

Chiral projection

$$j_{\mu}^{-} = \frac{-ig_W}{2\sqrt{2}} \bar{u} (1 + \gamma^5) \gamma^{\mu} (1 - \gamma^5) u = \frac{-ig_W}{2\sqrt{2}} \bar{u}_L \gamma^{\mu} u_L$$

Vector current interacting only with the left-handed particle, or right-handed anti-particle

# Weak interaction for neutrinos

- In case of neutrinos, both CC and NC interactions can be viewed as a vector current interacting only with the left-handed particle, or right-handed anti-particle
- Neutrinos only participate in weak interaction, they are produced in a left-handed eigen-state (chiral).
- If neutrinos are considered as no mass:
  - Chirality and helicity are the same
  - Neutrinos are always in left-handed state and no right-handed neutrinos exist!
- In fact neutrinos are massive, right-handed neutrino exists, e.g. as sterile neutrinos.

# Details of Neutrino Interactions

# Convenient variables used

- 4-momentum transfer:

$$Q^2 = -q^2$$

- Inelasticity:  $y = \frac{q \cdot P_{\text{target}}}{P_\nu \cdot P_{\text{target}}}$

In the target's rest frame:


$$y = \frac{q_0}{E_\nu} = \frac{E_\nu - E_l}{E_\nu}$$

- Bjorken scaling variable:

$$x = \frac{-q^2}{2P_{\text{target}} \cdot q}$$

- Invariant hadronic mass:

$$W = \sqrt{(q + P_{\text{target}})^2}$$



# Neutrino – Electron Interactions



# Neutrino – Electron interactions: CC

Threshold neutrino energy required:


$$E_\nu \geq m_l - m_e$$

Total cross-section in the center of mass frame  
(neglecting mass of electron and neutrino):

$$\sigma = \frac{1}{8\pi} \frac{g_W^4 E_\nu^2}{M_W^4} \left( 1 - \frac{m_\mu^2}{4E_\nu^2} \right)^2$$

Fermi's coupling:  $\frac{G_F}{\sqrt{2}} = \frac{g_W^2}{8M_W^2}$

Measurement in Charm II:  $\sigma(\nu_\mu e^-) = (1.651 \pm 0.093) \times 10^{-41} \left( \frac{E}{1 \text{ GeV}} \right) \text{ cm}^2$




# Neutrino – electron interactions: NC

- Incoming neutrinos can be:  $\nu_e$ ,  $\nu_\mu$  and  $\nu_\tau$
- There is no change in mass  $\rightarrow$  threshold-less interaction!
- Total cross-section (*9% of the inverse muon decay*):

$$\sin^2 \theta_W = 0.2324 \pm 0.0058 \pm 0.0059$$

$$\sigma_{TOT} = \frac{G_F^2 s}{\pi} \left( \frac{1}{4} - \sin^2 \theta_W + \frac{4}{3} \sin^4 \theta_W \right) = 1.4 \times 10^{-42} \text{ cm}^2 / \text{GeV} \cdot E_\nu (\text{GeV})$$

NC electron elastic scattering




# Neutrino – electron interactions: CC & NC

- In case of incoming neutrino  $\nu_e$   
 → In addition to NC, there is the second contributing, CC scattering.

→ The cross section is much bigger than the case of

$\nu_\mu$  and  $\nu_\tau$


$$\sigma(\nu_e e^-) = \frac{G_F^2 s}{\pi} \left[ \left( \frac{1}{2} + \sin^2 \theta_W \right)^2 + \frac{1}{3} \sin^4 \theta_W \right] = 0.96 \times 10^{-41} \left( \frac{E_\nu}{1 \text{ GeV}} \right) \text{ cm}^2$$




From

# Neutrino – electron interactions: Application

- To constraint neutrino flux using the uncertainty of cross-section




## Known Interaction (Standard Candle)



### Flux constraint using ND

$$\Phi = \frac{N}{\epsilon A \sigma}$$

Cross-section uncertainty goes into flux uncertainty



- $\nu$ -e scattering is well known interaction we can use to constrain the neutrino flux

**$\nu$ -e Scattering**

# Neutrino – Nucleons Interactions

# Neutrino – Nucleons interactions

1. Charged current quasi-elastic scattering:  $\nu_{\mu} + \mathbf{n} \rightarrow \mu^{-} + \mathbf{p}$
  2. Neutral current elastic scattering:  $\nu_{\mu} + \mathbf{N} \rightarrow \nu_{\mu} + \mathbf{N}$
  3. Single  $\pi, \eta, \mathbf{K}$  resonance productions:  $\nu_{\mu} + \mathbf{N} \rightarrow \mathbf{l} + \mathbf{N}' + \pi (\eta, \mathbf{K})$
  4. Coherent pion productions:  $\nu_{\mu} + \mathbf{X} \rightarrow \nu_{\mu} + \mathbf{X} + \pi_0$
  5. Deep inelastic scattering :  $\nu_{\mu} + \mathbf{N} \rightarrow \mathbf{l} + \mathbf{N}' + \mathbf{m}\pi(\eta, \mathbf{K})$
- $\mathbf{l}$ : lepton;  $\mathbf{N}, \mathbf{N}'$ : nuclons;  $\mathbf{m}$ : integer*

## Pion ( $\pi$ )

$$\pi^{+}(\bar{u}d); \pi^{-}(d\bar{u})$$

$$\pi^0(u\bar{u}/d\bar{d})$$

$$M_{\pi} \sim 140 \text{ MeV}$$

## Eta ( $\eta$ )

$$\eta = \frac{u\bar{u} + d\bar{d} - 2s\bar{s}}{\sqrt{6}}$$

$$M_{\eta} \sim 548 \text{ MeV}$$

## Kaon ( $\mathbf{K}$ )

$$\mathbf{K}^{+} = u\bar{s}; \mathbf{K}^{-} = s\bar{u}$$

$$\mathbf{K}^0 = d\bar{s}/s\bar{d}$$

$$m_{\mathbf{K}} \sim 495 \text{ MeV}$$

## Proton ( $\mathbf{p}$ )

$$\mathbf{p} = uud$$

$$m_{\mathbf{p}} \sim 940 \text{ MeV}$$


## Neutron ( $\mathbf{n}$ )

$$\mathbf{n} = udd$$


$$m_{\mathbf{n}} \sim 940 \text{ MeV}$$

# Neutrino – Nucleons interactions

- Nucleon target gives much more cross-section than electron target.
- **Elastic interactions:**
  - Dominate at small  $Q^2$
  - Nucleon recoil intact
  - CC interactions are referred to as “*quasi - elastic*” (change of charge and the mass transfer to the lepton in the final state).
- **Inelastic scattering:**
  - At low  $Q^2$ , resonance production is dominated.
  - At high  $Q^2$ , DIS production is dominated



# Neutrino – Nucleons interactions



## CC – $W^\pm$ exchange

- Quasi-elastic Scattering  
Target changes but no breakup  
 $\nu_\mu + n \rightarrow \mu^- + p$
- Coherent/Diffractive production  
Target unchanged  
 $\nu_\mu + n \rightarrow \mu^- + n + \pi^+$
- Nuclear resonance production  
Target goes to excited state and decays  
 $\nu_\mu + n \rightarrow \mu^- + p + \pi^0$  ( $N^*$  or  $\Delta$ )  
 $n + \pi^+$
- Deep Inelastic Scattering  
Target breaks up  
 $\nu_\mu + \text{quark} \rightarrow \mu^- + \text{quark}'$

## NC – $Z^0$ exchange

- Elastic Scattering  
Target unchanged  
 $\nu_\mu + n \rightarrow \nu_\mu + n$
- Coherent/Diffractive production  
Target unchanged  
 $\nu_\mu + N \rightarrow \nu_\mu + N + \pi^0$
- Nuclear resonance production  
Target goes to excited state and decays  
 $\nu_\mu + N \rightarrow \nu_\mu + N + \pi$  ( $N^*$  or  $\Delta$ )
- Deep Inelastic Scattering  
Target breaks up  
 $\nu_\mu + \text{quark} \rightarrow \nu_\mu + \text{quark}$

$q^2$



# NC Elastic and CC QE interactions


- In these processes of neutrino-nucleon interaction, a single lepton and a single nucleon are produced in which the nucleon recoils intact.
- In case of NC, for all flavours of neutrinos and anti-neutrinos, we have “*NC elastic*” scattering:



- In case of CC interaction, when neutrinos acquire sufficient energy:




→ Need energy to create the lepton’s mass → this is referred to as “*CC Quasi-elastic*” (CC QE) scattering.



# CC QE interactions


- CC QE interactions is an important channel for  $\nu$  oscillation experiments:
  - QE gives largest contribution to the cross-section of neutrino-nucleon interaction in a low region energy of neutrino ( $< 1$  GeV ).
  - QE is two body reaction  $\rightarrow$  the incident neutrino energy can be reconstructed from kinematics of the charged lepton  $\rightarrow$  for measuring oscillation parameters.



This implies that it is difficult to keep for the nucleon to remain intact at higher  $q^2$

# Resonance production


- This production obtained with higher  $Q^2$  transfer (or neutrino's energy,  $0.5 \text{ GeV} < E < 10 \text{ GeV}$ )  
→ inelastic scattering!
- The lepton part is almost the same as in the case of elastic scattering.
- In the hadronic part, the target nucleon is knocked into a baryon resonance → then decay into a nucleon + a single pion (mostly) or multi pions or Kaons or a radiative photon.





The available resonance depending on neutrino's energy

# Single pion production


Excitation of baryon resonance




Charged Current




Decay of baryon resonance



Neutral Current



# Single pion production



Main background of the nucleon decay:

Particles in the final state are the same as the ones from nucleon decay

Main background for the search of  $\nu_\mu \rightarrow \nu_e$  at T2K


In the NC scattering,  $\pi^0$  and  $\gamma$  production can mimic  $\nu_e$

Major contamination to the energy spectrum measurement


In the CC scattering,  $\pi$  production can be absorbed in the nucleus:

- $\pi$  can be considered as missing energy,  
 $\rightarrow$  background in searching for  $\nu_\mu \rightarrow \nu_\mu$  disappearance
- CC1pi can be mimicked as CCQE.

Charged Current




Neutral Current





# Single pion production cross-section




- **Data points:** total inclusive  $\nu_\mu$  CC interaction  $\nu_\mu p \rightarrow \mu^- p \pi^+$

$\nu_\mu$  CC 1pi cross-section on deuterium

This process is dominated in the range  $0.5 \text{ GeV} < E < 10 \text{ GeV}$

# Coherent Interactions (nuclear target)



- Pion production without breaking the target nucleus.
- Cross-section is smaller than the resonance-mediated production.
- At low range of  $E_\nu$  : NC scattering:  
$$\nu + X \rightarrow \nu + X + \pi_0$$
- Recently, cross-section of **charged current** coherent pion production ( $\nu + {}^{12}\text{C} \rightarrow l^\pm + {}^{12}\text{C} + \pi^0$ ) was found to be very small in  $\sim < \text{GeV}$  region.
- CC experimentally observed in higher  $E_\nu$






# Deep Inelastic Scattering

- DIS process appears from  $E > \text{few GeV}$
- Nucleons are made of quarks.
- Understood as neutrino – quark interaction.
- $E_\nu$  is calculated as energy of lepton + energy of hadrons.



Hadrons

# Deep Inelastic Scattering



• **Data points:** total inclusive  $\nu_\mu$  CC interaction measured on different targets.

$\nu_\mu$  CC DIS cross-section on deuterium

DIS process dominates for  $E_\nu > 10$  GeV and increases linearly with  $E_\nu$  until W, Z mass!


# Neutrino – Nucleus Interactions

# Neutrino – nucleus interactions: motivation

- Limitation of **neutrino – electron interaction**:
  - Well-understood but cross-section is much smaller than neutrino – nucleon interactions
  - Impossible to create a target made of only free electrons in the reality.
- Experiments think of **neutrino – nucleons interaction**:
  - Impossible to construct a target made of pure neutrons
  - Hydrogen target containing only proton → CC QE interaction is only for anti-neutrinos with lower cross-section.
  - Deuterium made of proton and neutron is a good target but it is light → low interaction rate.
- Neutrino-nucleus interactions:
  - Give higher interaction rate
  - In experiments, detectors are build using heavier nuclei such as carbon, oxygen or iron
  - Nuclear effects are present making complication in understanding interactions observed in detectors!







# Neutrino-nucleus interactions: nuclear effects

- Initial state of the nucleons:
  - Nucleons in a nucleus move around inside the nuclear potential, changing their momentum and direction.
  - The direction and momentum of the nucleon affects the kinematics of any interaction
  - The initial momentum spectra of nucleons is not well known.
  - → Need models to describe this.
- Final State Interaction (FSI)
- Nuclear effects become more important at low energy region.
- Nuclear effects are simulated by MC generators.



The nucleon momentum distributions from a RFG and a SF (both for Carbon)

# Lepton kinematics



# Introduction to Neutrino Event Generators


# Tasks of Neutrino event generators

- **Simulate neutrino interactions:** each generator is expected to simulate all possible interactions using appropriate models.
- **Simulate signals and backgrounds observed in the detector** → background can be extracted from real data.
- **A bridge/tool to compare real data and theories** in order to extract neutrino oscillation parameters.
- **Reduce systematic uncertainties in measuring physics observation** by reducing uncertainties caused by the understanding of neutrino interactions with nucleus.
  - precision of the neutrino event generators is required to better understand neutrino interactions.
- **Can be used to evaluate systematic uncertainties in extracting the physics results.**



# Neutrino event generators

- What? → Neutrino event generators are softwares simulating neutrino interactions.
- How do they work?



# Examples of Neutrino Event Generators

## NEUT

- Developed initially for Kamiokande exp. then for Super-K, K2K, SciBooNE and T2K.
- Used for Super-K and T2K MC official production
- Mainly written in Fortran

## GENIE


- Developed by an international collaboration
- Universal neutrino event generator
- Written in C++ and well maintained, open source

## NuWro

- More theory oriented. Developed by people from Wroclaw University
- Written in C++.

# Challenges with Neutrino Scattering

- $\nu$  beams are not mono-energetic:
  - $\nu$  flux
  - At a given  $E_\nu$ , there is contributions from multi processes
- $\nu$  cross-section is not well-constrained in a region interested.
- Targets used in  $\nu$  experiments are nucleus  $\rightarrow$  nuclear effects are included in cross-section and kinematics of out-going particles:
  - $\rightarrow$  MC generators need to be included these effects.



# Neutrino flux and cross-section

- Frequency of neutrino interactions when the flux is uniform:

$$f = \phi \left[ \frac{1}{\text{time} * \text{area}^2} \right] * N * \sigma[\text{area}^2]$$

Flux                      Number of scattering centers                      Cross-section


- In reality,  $\mathcal{V}$  flux is non-monoenergetic:

$$f = \int dE_{\nu} \phi(E_{\nu}) * N * \sigma(E_{\nu})$$


- Neutrino event generators are used to **predict flux**, to **simulate detector response** and to **generate interactions**.

# Example of Neutrino flux and Event rate

Flux



Event Rate = Flux x Cross-section



# Introduction to NEUT

- NEUT is a program simulating neutrino interactions with  $E_\nu \in (100 \text{ MeV}, \sim \text{TeV})$
- Target nucleus used: primarily proton and Oxygen then also rather well tested with Carbon ( at K2K, SciBooNE )

## Works carried out by NEUT:

- Provides cross-sections to estimate the interaction rate or to select the interaction mode.
- Simulates primary neutrino interaction with nucleon and nucleus targets.
- Simulates meson interactions in the target, especially in detail for the low momentum pion.
- Simulates nucleon re-scattering in the target nucleus.

Thank you for you attention!