

# Self-introduction




My name is Atsumu Suzuki  
(鈴木州) from Kobe  
University  
Member of SK, HK, and T2K



Scuba diving  
(a couple of times a year)



Tennis  
weekend



My shoosie

Skating in winter



# T2K experiment

1. Introduction
2. Experimental setup
  - 2.1 Experimental overview
  - 2.2 Accelerator
  - 2.3 Beam line
  - 2.4 Detectors
3. Current status & results
4. Future prospect

Super-K

J-PARC

Image NASA  
© 2007 Europa Technologies  
Image © 2007 TerraMetrics  
© 2007 ZENRIN

© 2007 Google™

# 1. Introduction



# Neutrino oscillation measurement before T2K



SK, SNO, KamLand

$(\theta_{12} \doteq 34^\circ$   
 $\Delta m_{21}^2 \sim 8 \times 10^{-5} \text{ eV}^2)$

$\Delta m_{ij}^2 = m_i^2 - m_j^2$


Not observed

$\theta_{13} = ?$ ,  $\delta_{CP} = ?$



SK, K2K, MINOS

$\theta_{23} \doteq 45^\circ$   
 $\Delta m_{32}^2 \sim 2.5 \times 10^{-3} \text{ eV}^2$



Which mass hierarchy ,  
 $m_1 < m_2 < m_3$  (Normal Hierarchy, NH) or  
 $m_3 < m_1 < m_2$  (Inverted Hierarchy, IH) ?



# T2K experiment



- **Tokai to Kamioka (T2K) long-baseline neutrino oscillation experiment**
- **Muon neutrinos from J-PARC  $\Rightarrow$  Super-Kamiokande @ 295 km**
- **T2K accomplishments**
  - **Discovery of  $\nu_e$  appearance in 2013**
  - **World-best precision measurement of  $\nu_\mu$  disappearance**

# The main goals of T2K

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta_{CP}} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta_{CP}} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$

Flavor  
eigen states

Atmospheric term

Reactor term

Solar term

Mass  
eigen states

$$C_{ij} = \cos \theta_{ij}, S_{ij} = \sin \theta_{ij}$$


- $\nu_\mu$  disappearance (We measure how much  $\nu_\mu$ 's disappear after the flight.)  
 $\Rightarrow \theta_{23}, \Delta m_{32}^2$  precision measurement (Neutrino energy of T2K is low and  $\tau$  cannot be produced.\*)
- $\nu_e$  appearance (We measure how much  $\nu_e$ 's which do not exist originally appear after the flight.)  
 $\Rightarrow \theta_{13}, \delta_{CP}$  measurement

\*Homework 1

How much energy do we need for a neutrino to produce a  $\tau$ ?



# T2K $\delta_{CP}$ measurement



For  $\delta_{CP}$ , look for  $\nu/\bar{\nu}$   
difference of  $\nu_e$  appearance

$$P(\nu_\mu \rightarrow \nu_e) = P(\bar{\nu}_\mu \rightarrow \bar{\nu}_e) ?$$

# T2K Collaboration




~500 members, 76 institutes, 13 countries (+CERN)





## **2. Experimental setup**


# 2.1 Experimental overview



- **High intensity proton beam hits the graphite target.**
- Secondary  $\pi/K$ 's focused by magnetic horns and decay to neutrinos
- Neutrino beam from  $\pi^+ \rightarrow \mu^+ + \nu_\mu$
- Antineutrino beam from  $\pi^- \rightarrow \mu^- + \bar{\nu}_\mu$
- Changing neutrino beam mode by flipping the horn polarity
- Beam dump absorbs remained hadrons & the direction & intensity of the muons from  $\pi(K) \rightarrow \mu + \nu$  are measured by MUMON.
- Unoscillated neutrinos are measured by near detectors
- Neutrinos are measured by far detector 295km far away from target




# 2.1 Experimental overview



- High intensity proton beam hit the graphite target
- **Secondary  $\pi/K$ 's focused by magnetic horns and decay to neutrinos**
- Neutrino beam from  $\pi^+ \rightarrow \mu^+ + \nu_\mu$
- Antineutrino beam from  $\pi^- \rightarrow \mu^- + \bar{\nu}_\mu$
- **Changing neutrino beam mode by flipping the horn polarity**
- Beam dump absorbs remained hadrons & the direction & intensity of the muons from  $\pi(K) \rightarrow \mu + \nu$  are measured by MUMON.
- Unoscillated are neutrinos measured by near detectors
- Neutrinos are measured by far detector 295km far away from target


# 2.1 Experimental overview



- High intensity proton beam hit the graphite target
- Secondary  $\pi/K$ 's focused by magnetic horns and decay to neutrinos
- Neutrino beam from  $\pi^+ \rightarrow \mu^+ + \nu_\mu$
- Antineutrino beam from  $\pi^- \rightarrow \mu^- + \bar{\nu}_\mu$
- Changing neutrino beam mode by flipping the horn polarity
- Beam dump absorbs remained hadrons, and the direction & intensity of the muons from  $\pi(K) \rightarrow \mu + \nu$  are measured by MUMON.
- Unoscillated neutrinos are measured by near detectors
- Neutrinos are measured by far detector 295km far away from target




# 2.1 Experimental overview



- High intensity proton beam hit the graphite target
- Secondary  $\pi/K$ 's focused by magnetic horns and decay to neutrinos
- Neutrino beam from  $\pi^+ \rightarrow \mu^+ + \nu_\mu$
- Antineutrino beam from  $\pi^- \rightarrow \mu^- + \bar{\nu}_\mu$
- Changing neutrino beam mode by flipping the horn polarity
- Beam dump absorbs remained hadrons & the direction & intensity of the muons from  $\pi(K) \rightarrow \mu + \nu$  are measured by MUMON.
- **Unoscillated neutrinos are measured by near detectors.**
- Neutrinos are measured by far detector 295km far away from target.


# 2.1 Experimental overview




- High intensity proton beam hit the graphite target
- Secondary  $\pi/K$ 's focused by magnetic horns and decay to neutrinos
  - Neutrino beam from  $\pi^+ \rightarrow \mu^+ + \nu_\mu$
  - Antineutrino beam from  $\pi^- \rightarrow \mu^- + \bar{\nu}_\mu$
  - Changing neutrino beam mode by flipping the horn polarity
- Beam dump absorbs remained hadrons & the direction & intensity of the muons from  $\pi(K) \rightarrow \mu + \nu$  are measured by MUMON.
- Unoscillated neutrinos are measured by near detectors
- Neutrinos are measured by far detector (SK) 295km far away from target.



# 2.1 Experimental overview




- **2.5° off-axis neutrino beam**
  - Narrow band beam below 1 GeV
  - Peak at 0.6 GeV at oscillation max w/ 295km
  - Small high energy tail
  - High energy neutrino background can be suppressed





# 2.2 J-PARC

## (Japan Proton Accelerator Research Complex)





## Linac (Linear accelerator)

- The first stage accelerator, **330 m** in length.
- Protons are accelerated to **400 MeV**.



## RCS

### (Rapid Cycling Synchrotron)

- The second stage accelerator, Proton Synchrotron of **348 m** circumference.
- Protons are accelerated up to **3 GeV**.




## Main Ring

- The third (and final) stage accelerator. Proton Synchrotron of **1568 m** circumference.
- The **30 GeV** proton beam is extracted to the neutrino beamline and to hadron hall.



## Proton beam to neutrino beam line





**Bunch**

1 spill (= 8 bunches)

## **2.3 Beam line**



# Primary Proton beam line





Primary Beam Monitor	Number
I : Beam Intensity (CT)	5
C: Beam Position (ESM)	21
P: Beam Profile (SSEM)	19
P: Beam Profile (OTR)	1
Beam Loss Monitor	50





Normal-conducting magnet in Preparation section

# Beam monitors



$$x \propto \frac{CH0 - CH1}{CH0 + CH1}$$


$$y \propto \frac{CH2 - CH3}{CH2 + CH3}$$




**Secondary Segmented Emission Monitor (SSEM)**

# Optical Transition Radiation monitor (OTR)


Transition radiation is a form of EM radiation emitted when a charged particle crosses the boundary between two media with different permittivities.



$$\frac{dN_{\text{photon}}}{d\lambda} = \frac{2\alpha}{\pi\lambda} \left\{ \ln \left( \frac{\gamma\lambda}{\lambda_{pe}} \right) - 1 \right\}$$



F. Sakamoto, et. al - Emission and energy measurements of low-energy electrons beam using optical transition radiation techniques, JJAP vol.44, 3, 2005, 1485-1491.



**OTR online event display**

OTR Image Display

Command Menu: Pause/Resume, Reset History, Save Current Image, Exit

OTR History Images | OTR Current Event | Position History | Sigma History

2009-04-24 20:32:50 UTC Spill: 56672

Distance Corrected Image

X Projection - Spill 56672




Position = -1.56 mm  
Sigma = 2.84 mm

Y Projection - Spill 56672


Position = -0.92 mm  
Sigma = 1.87 mm




# Target Station



# How to produce neutrino beam




We have to make the 2ry charged particle direction parallel and inject neutrinos to the target direction.



**Magnetic horn !**



# Magnetic horn





# Magnetic horn

- Magnetic horn is **focus/defocus** device installed just downstream of the target. High current (320 kA) generate magnetic field, and direction  $\pi$ 's are changed.
- In  $\nu$ -mode,  $\pi^+$  is focused and  $\pi^-$  is defocused. Accordingly,  $\nu$ -rich beam is generated in the forward direction.
- In  $\bar{\nu}$ -mode,  $\pi^-$  is focused, and  $\pi^+$  is defocused.  $\bar{\nu}$ -rich beam is generated in the forward direction.
- Neutrino flux at SK is enhanced by factor  **$\sim 10$**  (total) and  **$\sim 16$**  (at  $\sim 0.6\text{GeV}$ ).

- Neutrino components in each mode:

	$\nu_\mu$	$\bar{\nu}_\mu$	$\nu_e + \bar{\nu}_e$
$\nu$ -mode	$\sim 97\%$	$\sim 2\%$	$\sim 1\%$
$\bar{\nu}$ -mode	$\sim 2\%$	$\sim 97\%$	$\sim 1\%$






# Decay Volume and Beam Dump

## Muon monitors in Muon Pit



Remaining protons and undecayed  $\pi$ -mesons are absorbed by a **beam dump** composed of large graphite blocks.




neutrino beamline

## Beam Dump (installation)



Installation of decay volume



Inside of decay volume (rectangular shape)



# Muon monitors

- Two types of muon monitors are installed downstream of the beam dump for redundancy.


## Ionization chamber

7x7 channels  
Ar+2%N<sub>2</sub> gas (~Mar. 2015)  
He+1%N<sub>2</sub> gas (May~ 2015)



## Semiconductor array

7 x 7 channels of Silicon PIN photo diode.




- Confirm the position of the beam center with **< 3cm** resolution on a bunch by bunch basis. This corresponds to **< 0.3mrad** beam direction accuracy.

## **2.4 Detectors**

# Near Detectors at 280m downstream

- The detectors were made in the underground experimental hall, **33.5m** depth and **17.5m** diameter. It is located at **280m** downstream from the target.
- Two detectors were installed; they are **On-axis Detector** in the direction of the neutrino beam center, and **Off-axis detector** in the direction of Super-Kamiokande.




Unlike K2K, a water Cherenkov detector cannot be used in T2K near detector.

- Event rate is too high.
- Neutrino energy is high and muons escape from an 1kt tank.




# On-axis detector (INGRID)


- Consists of **16** modules; **7** horizontal, **7** vertical, and **2** off-diagonal. Each module is 1m x 1m x 1m cube.
- Each module is “sandwich” of **11 plastic scintillator** layers and **10 iron** layers. They are surrounded by **4** veto planes.
- The neutrino beam center is obtained from horizontal/vertical distributions of the neutrino event rate. The nominal accuracy is **~0.1 mrad**.




INGRID event view



Number of event distribution




# Stability of event rate and beam direction



- Event rate is stable over neutrino and anti-neutrino periods.
- Beam direction is much stable than our requirement, **1 mrad**.



# ND280 detector





neutrino beam





# Off-axis detector (ND280)

- **ND280** is made of several components.
- 2 **FGDs** (Fine-Grained Detectors) consist of scintillator bars. FGD2 has water as a target material.
- 3 gas-filled **TPCs** (Time Projection Chambers) track charged particles.
- All components are in **0.2 T** of magnetic field. The **magnets** were previously used in UA1\* and NOMAD.
- Charged particles are deflected by the magnetic field. The curvature of the track recorded by TPC are used to determine the momentum of the particles.
- Neutrino flux as well as neutrino interactions can be studied from the reconstructed track information.
- Other components are **P0D** ( $\pi^0$  detector), **ECAL**(Electromagnetic CALorimeter) and **SMRD**(Side Muon Range Detector).





\* W & Z were discovered by this experiment.

# Magnet yolk






# Magnet yolk installation






# SMRD scintillator

width:  
167 mm for horizontal gaps  
175 mm for vertical gaps




Wave length shifting fiber

Scintillator thickness: 7.1 mm  
Groove depth: 2.5 mm




Installed in each gap



4 or 5 scintillators  
are combined  
into one module.



# Easy exercise for a range detector




Assume that a  $\mu^-$  stops in the mid. of the Fe layer  
Densities of Fe & the scintillator : 8 g/cm<sup>3</sup> & 1 g/cm<sup>3</sup>, respectively.  
 $dE/dx \sim 2 \text{ MeV}/(\text{g}/\text{cm}^2)$

Calculate the muon deposit energy:  $E_{\mu}^{\text{deposit}}$

$$\begin{aligned} E_{\mu}^{\text{deposit}} &= (5 \text{ cm}/\sin 30^{\circ} \times 8 \text{ g}/\text{cm}^3 + 2 \text{ cm}/\sin 30^{\circ} \times 1 \text{ g}/\text{cm}^3) \times 2 \text{ MeV}/(\text{g}/\text{cm}^2) \\ &= \mathbf{168 \text{ MeV}} \end{aligned}$$



# WAGASCI & BabyMIND detectors (1.5 degrees off-axis)



- Water filled plastic scintillator Lattice and magnetised tracking (BabyNIND) detectors
- We plan to use in the oscillation analysis
- First WAGASCI cross section paper: [Phys. Rev. D 97, 012001](#)




# Near Detector Complex





# Far detector : Super-Kamiokande (SK)

- 50kt water Cherenkov detector. The fiducial volume of the inner detector is **22.5** kton, and is viewed by **11129** 20-inch diameter PMTs. Outer water layer surrounding the inner volume is viewed by 1885 8-inch diameter PMTs.
- Located at 1000 m underground in Kamioka mine, Japan. The distance from the J-PARC is **295** km.
- Now Gd was loaded for enhanced neutron detection to separate between  $\nu_e$  and  $\bar{\nu}_e$  interactions.



# **3. Current status & results**



# Parameters related to beam power

- **Beam power**

$$\text{Energy of one proton} \times \text{Protons per second (pps)}$$

Let's calculate yourself for T2K case.

- Proton energy: 30 GeV,  
Protons per cycle:  $2.67 \times 10^{14}$ ,  
Cycle time: 2.48 sec  
(1 GeV =  $10^9$  eV, 1 eV =  $1.6 \times 10^{-19}$  J)
- Answer in "kW".

$$30 \text{ GeV} \times 2.67 \times 10^{14} / 2.48 \times 1.6 \times 10^{-19} \times 10^{-3} \text{ kW} = 517 \text{ kW}$$

K2K proton energy: 12 GeV, protons per cycle:  $6 \times 10^{12}$  cycle time: 2.2 sec, beam power: 5 kW


- **POT (protons On Target)**

$$\text{pps} \times \text{Beam time}$$

\*Homework 2

How many years do we need to get  $1 \times 10^{22}$  POT if the beam power is 517 kW as described above ?

# T2K Data Accumulation Summary



**Total POT for the results here:  $3.60 \times 10^{21}$  ν mode:  $1.97 \times 10^{21}$  POT**  
**(Protons on Target) ν̄ mode:  $1.63 \times 10^{21}$  POT**



(Run11,  $1.78 \times 10^{20}$  POT in ν mode, the very first T2K data with a Gd-loaded SK)

**766 kW eq. pulse acceleration was achieved in 1 shot operation on Apr. 16, 2023!!**

# Event Selection at the Far Detector (SK)

## Event Selection Criteria

1. Total energy deposit in the inner detector is larger than **30 MeV** equivalent.
2. No outer detector activity
3. The event time agrees with  **$\sim 5 \mu\text{sec}$**  beam period in **2.48 sec** accelerator cycle.  
(8 bunch structure can be found.)
4. 1 Ring events  
→  $e/\mu$  particle identification is applied






# $\mu/e$ identification in Super-Kamiokande


$$\nu_{\mu} \rightarrow \mu$$

Only direct Cherenkov light from  $\mu$   
Clear Cherenkov ring edge




$$\nu_e \rightarrow e$$

Cherenkov light from e-m shower.  
Electrons and positrons are heavily scattered.  
Cherenkov ring edge is fuzzy.



- $\mu/e$  misidentification probability is less than **1 %**.




# Event Selection at the Far Detector (cont'd)

Examine Particle ID of 1 ring events

## $\nu_\mu$ selection


- $\mu$ -like PID
- $p_\mu > 200 \text{ MeV}/c$
- Michel electron 1 or 0



## $\nu_e$ selection

- e-like PID
- $p_e > 100 \text{ MeV}/c$
- Michel electron 0
- $E_{\text{rec}} < 1250 \text{ MeV}$
- $\pi^0$  rejection


$\pi^0$  rejection :  
Forced 2<sup>nd</sup> ring is  
assumed. Invariant  
mass and likelihood  
for  $\pi^0$  are examined.





# The first neutrino event in the T2K experiment

Recorded on  
**Feb-24-2010**






# $\nu_\mu$ disappearance analysis

CCQE (Charged Current Quasi-Elastic)


$\nu_\mu + n \rightarrow \mu^- + p$  candidate



## Disappearance of $\nu_\mu/\bar{\nu}_\mu$ :


$$\begin{aligned}
 P(\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu) &= 1 - 4\cos^2(\theta_{13})\sin^2(\theta_{23}) \\
 &\times \left[ 1 - \cos^2(\theta_{13})\sin^2(\theta_{23}) \right] \sin^2\left(\frac{\Delta m_{32}^2 L}{4E}\right) \\
 &+ (\text{solar, matter effect terms})
 \end{aligned}$$

location of dip:  $\Delta m_{32}^2$   
 depth of dip:  $\sin^2(\theta_{23})$




# Recent result of $\nu_{\mu}/\bar{\nu}_{\mu}$ (1)

T2K Run 1-10 Preliminary






T2K Run 1-10 Preliminary







# Far detector new samples

- $\nu_\mu$  CC  $1\pi^+$  interactions in  $\nu$ -mode were added to CCQE events.
- “Two rings ( $1\mu^-$  and  $1\pi^+$ ) and Michel electron (from  $1\mu^-$ )” or “one  $1\mu^-$  ring and 2 Michel electrons (from  $1\mu^-$  &  $1\pi^+$ )”
- Increase  $\nu$ -mode  $\mu$ -like statistics by  $\sim 30\%$



# Measurements on $\theta_{23}$ and $\Delta m_{32}^2$



		$\sin^2 \theta_{23}$		Sum
		$< 0.5$	$> 0.5$	
$\Delta m_{32}^2$	$> 0$ (NO)	0.195 (0.260)	0.613 (0.387)	0.808 (0.647)
	$< 0$ (IO)	0.035 (0.152)	0.157 (0.201)	0.192 (0.353)
Sum		0.230 (0.412)	0.770 (0.588)	1.000

Fractions of posterior probability in different combinations of the mass ordering and  $\theta_{23}$  octant from fit to T2K data with (without) the reactor constraint on  $\sin^2 \theta_{13}$ .

arXiv:2303.05222 (to be appeared in Eur. Phys. J. C)

Slight preference for **upper octant** ( $\sin^2 \theta_{23} > 0.5$ ) and **normal mass ordering**.

## Appearance of $\nu_e/\bar{\nu}_e$ :


$$P(\bar{\nu}_\mu \rightarrow \bar{\nu}_e) =$$

$$\begin{aligned} & \sin^2(\theta_{23}) \sin^2(2\theta_{13}) \sin^2\left(\frac{\Delta m_{32}^2 L}{4E}\right) \\ & -^{(+)} \left[ \sin(2\theta_{12}) \sin^2(2\theta_{23}) \sin^2(2\theta_{13}) \cos(\theta_{13}) \right. \\ & \times \sin\left(\frac{\Delta m_{21}^2 L}{4E}\right) \sin^2\left(\frac{\Delta m_{32}^2 L}{4E}\right) \sin(\delta_{CP}) \\ & \left. + (\text{CP-even, solar, matter effect terms}) \right] \end{aligned}$$

magnitude of the peak:

$$\sin^2(\theta_{23}), \sin^2(2\theta_{13}), \delta_{CP}$$

$$P(\nu_\mu \rightarrow \nu_e) = P(\bar{\nu}_\mu \rightarrow \bar{\nu}_e) \text{ or not?}$$







# T2K Results Restrict Possible Values of Neutrino CP Phase



The T2K Collaboration has published new results showing the strongest constraint yet on the parameter that governs the breaking of the symmetry between matter and antimatter in neutrino oscillations. Using beams of muon neutrinos and muon antineutrinos, T2K has studied how these particles and antiparticles transition into electron neutrinos and electron antineutrinos, respectively. The parameter governing the matter/antimatter symmetry breaking in neutrino oscillation, called  $\delta_{cp}$  phase, can take a value from  $-180^\circ$  to  $180^\circ$ . For the first time, T2K has disfavored almost half of the possible values at the 99.7% ( $3\sigma$ ) confidence level, and is starting to reveal a basic property of neutrinos that has not been measured until now. This is an important step on the way to knowing whether or not neutrinos and antineutrinos behave differently. These results, using data collected through 2018, have been published in the multidisciplinary scientific journal, Nature on April 16. ([Nature | Vol 580 | 16 April 2020 |](#))

# Recent result of $\nu_e/\bar{\nu}_e$

T2K Run 1-10, 2022 Preliminary



Parameter	With reactor constraint	
	Normal ordering	Inverted ordering
$\delta_{CP}$ (rad.)	$-1.97^{+0.97}_{-0.62}$	$-1.44^{+0.56}_{-0.59}$
$\sin^2 \theta_{13}/10^{-3}$	—	—
$\sin^2 \theta_{23}$	$0.561^{+0.019}_{-0.038}$	$0.563^{+0.017}_{-0.032}$
$\Delta m_{32}^2/10^{-3}$ (eV <sup>2</sup> )	$2.494^{+0.041}_{-0.058}$	—
$ \Delta m_{31}^2 /10^{-3}$ (eV <sup>2</sup> )	—	$2.463^{+0.042}_{-0.056}$

# **4. Future prospect**



# Future extension

- Upgrades

- Beam :MR power supply → > 800 kW by 2023


- MR RF upgrade → > 1 MW by 2027

- ND280: Super FGD, HA-TPC, & TOF (→ next page)

- Aiming for >  $3\sigma$  sensitivity for CP violation with significantly improved statistics


In April 2023  
Successful demonstration of  
MR-FX 30 GeV acceleration  
766 kW eq. ( $2.17 \times 10^{14}$  ppp) in 1.36 s cycle

MR upgrade (history & plan)



# Near Detector Upgrade

arXiv:1901.03750







P0D (to measure NC  $\pi^0$  production) was replaced by a new scintillator target (Super FGD), 2 High Angle TPCs, and TOF.





# New detectors

## Super-FGD



- \* New concept of detectors,  $2 \times 10^6$  1cm<sup>3</sup> cubes
- \* Each cube is read by 3 WLS → 3D view

## High-Angle TPCs



- \* New TPCs instrumented with Encapsulated Resistive Anode MicroMegas (ERAM)

## TOF



- 6 TOF planes to reconstruct track direction
- Time resolution ~150 ps



# Super FGD

(vii) Horizontal fibers assembly




(viii) Wall MPPCs assembly



(ix) Vertical fibers assembly






(x) Top MPPCs assembly



(xi) LED calib. modules assembly



(xii) Light barrier/cables assembly



**Super-FGD assembly has been completed at J-PARC in April.**

**First cosmic ray tracks have been observed.**



ICISE

INTERNATIONAL CENTRE FOR INTERDISCIPLINARY SCIENCE AND EDUCATION

TRUNG TÂM QUỐC TẾ KHOA HỌC VÀ GIÁO DỤC LIÊN NGÀNH  
CENTRE INTERNATIONAL DE SCIENCE ET D'ÉDUCATION INTERDISCIPLINAIRES  
QUY NHƠN - VIET NAM

Thank you for listening (^o^)



ICISE

INTERNATIONAL CENTRE FOR INTERDISCIPLINARY SCIENCE AND EDUCATION

TRUNG TÂM QUỐC TẾ KHOA HỌC VÀ GIÁO DỤC LIÊN NGÀNH  
CENTRE INTERNATIONAL DE SCIENCE ET D'ÉDUCATION INTERDISCIPLINAIRES  
QUY NHƠN - VIET NAM

Thank you for listening (^o^)

\*Homework 3  
Who is he ?





# Solutions of the homeworks

## \*1 How much energy do we need for a neutrino to produce a $\tau$ ?

(Answer) When a  $\tau$  particle is produced by CCQE interaction, i.e.

$$\nu_\tau + n \rightarrow \tau^- + p,$$

the center of mass energy squared  $s$  is

$$s = (E_\nu + m_n)^2 - p_\nu^2 = m_n^2 + 2m_n E_\nu,$$

since  $E_\nu \approx p_\nu$ . The threshold condition is

$$s = (m_\tau + m_p)^2 = m_n^2 + 2m_n E_\nu.$$

Hence the threshold energy is

$$E_\nu = \frac{(m_\tau + m_p)^2 - m_n^2}{2m_n} = \frac{(1.777 + 0.9383)^2 - 0.9396^2}{2 \times 0.9396} = \underline{\underline{3.45 \text{ [GeV]}}}$$

## \*2 How many years do we need to get $1 \times 10^{22}$ POT ?

(Answer) Beam time = POT/pps =  $1 \times 10^{22} / 2.67 \times 10^{14} / 2.48 = 9.29 \times 10^8$  [sec]

$$= \underline{\underline{29.5 \text{ [yr]}}}$$

(This number is not realistic and we cannot operate the accelerator whole year.

So we need much more beam power actually.)

## \*3 Who is he ? (Answer) **Of course, Son Cao-san !!**